

PENGARUH LATIHAN TAICHI UNTUK RESIKO JATUH PADA LANSIA DI UPT PSTW BANYUWANGI

EVIDANCE BASED NURSING



Disusun Oleh:

Ade Nur Alfa Rihidiansyah	21101001
Anisatul Islamiyah	21101007
Efiq Elvira Rismasita	21101019
Karina Maya Ovie Andariesta	21101047
Mitha Anggraini	21101060
Renno Indra Nugroho	21101000
Wahyu Triya Kusuma	21101102

**PROGRAM STUDI PROFESI NERS
FAKULTAS ILMU KESEHATAN
UNIVERSITAS dr. SOEBANDI
2022**

LEMBAR PENGESAHAN

Evidence Based Nursing yang berjudul "*Pengaruh Senam Tai Chi Terhadap Kejadian Resiko Jatuh Pada Lansia di Wisma Sri Tanjung UPT PSTW Banyuwangi*" telah diperiksa dan disahkan pada :

Hari : Kamis

Tanggal : 31 Maret 2022

Yang Mengesahkan,

Pembimbing Klinik
UPT PSTW Banyuwangi




Diana Efendi, Amd. Kep

Pembimbing Akademik
Universitas dr. Soebandi Jember



Irwina Angelia Silvanasari, S.Kep., Ns., M.Kep
NIDN. 0709099005

Mengetahui,
Kepala UPT PSTW Banyuwangi



Drs. Agung Pambudi, M.Si
NIP. 19670814 19930 1 011

HALAMAN PERSEMBAHAN

Segala puji bagi Allah SWT atas limpahan rahmat dan Ridho-nya yang senantiasa selalu memberikan kemudahan, petunjuk, kekuasaan, dan keyakinan sehingga saya dapat menyelesaikan penyusunan EBN ini tepat pada waktunya.

DAFTAR ISI

HALAMAN PERSEMBAHAN	ii
MOTTO	Error! Bookmark not defined.
HALAMAN PERNYATAAN ORISINALITAS...	Error! Bookmark not defined.
LEMBAR PERSETUJUAN.....	Error! Bookmark not defined.
HALAMAN PENGESAHAN.....	Error! Bookmark not defined.
ABSTRAK	Error! Bookmark not defined.
KATA PENGANTAR	Error! Bookmark not defined.
DAFTAR ISI.....	iv
DAFTAR TABEL	vii
DAFTAR GAMBAR	viii
BAB 1 PENDAHULUAN	1
1.1 Latar Belakang	1
1.2 Rumusan Masalah	5
1.3 Tujuan penelitian	5
1.3.1 Tujuan Umum	5
1.3.2 Tujuan Khusus	5
1.2 Manfaat Penelitian	6
1.4.1 Peneliti	6
1.4.2 Institusi.....	6
1.4.3 Masyarakat.....	Error! Bookmark not defined.
BAB 2 TINJAUAN PUSTAKA	7
2.1 Konsep lansia	7
2.1.1 Definisi lansia	7
2.1.2 Batasan Lanjut Usia	8
2.2 Konsep Resiko jatuh Tubuh	9

2.2.1	Perubahan Resiko jatuh Tubuh Dalam Proses Menua.....	9
2.2.2	Komponen Resiko jatuh Tubuh	10
2.2.3	Penilaian Fungsi Resiko jatuh.....	10
2.2.4	Faktor-Faktor yang Mempengaruhi Resiko jatuh Tubuh Lansia.....	10
2.3	Konsep Senam Taichi.....	12
2.3.1	Pengertian Senam Taichi.....	12
2.3.2	Tujuan senam taichi	12
2.3.3	Prinsip Senam Taichi	13
2.3.4	Penatalaksanaan Senam Taichi	15
2.4	Kerangka Teori.....	Error! Bookmark not defined.
BAB 3 METODE PENELITIAN.....		Error! Bookmark not defined.
3.1	Strategi Pencarian Literature	Error! Bookmark not defined.
3.1.1	Protokol dan Registrasi Desain penelitian.....	Error! Bookmark not defined.
3.1.2	Database Pencarian	Error! Bookmark not defined.
3.1.3	Kata Kunci	Error! Bookmark not defined.
3.2	Kriteria Inklusi dan Eksklusi.....	Error! Bookmark not defined.
3.3	Seleksi Studi dan Penilaian Kualitas ...	Error! Bookmark not defined.
3.3.1	PRISMA <i>Flow Diagram</i>	Error! Bookmark not defined.
3.2.2	Hasil Pencarian <i>Literature Review</i> ...	Error! Bookmark not defined.
BAB HASIL DAN ANALISA		Error! Bookmark not defined.
4.1	Hasil	Error! Bookmark not defined.
4.1.1	Hasil Pencarian Literatur.....	Error! Bookmark not defined.
4.1.2	Karakteristik Studi.....	Error! Bookmark not defined.
4.1.3	Karakteristik Responden Studi.....	Error! Bookmark not defined.
4.2	Analisis	Error! Bookmark not defined.
4.2.2	Latihan Taichi Untuk Kemampuan Resiko jatuh Pada Lanjut Usia	Error! Bookmark not defined.

BAB 5 PEMBAHASAN	34
5.1 Kemampuan Resiko jatuh pada Lanjut Usia.....	36
5.2 Latihan Taichi terhadap Kemampuan Resiko jatuh pada Lanjut Usia.....	37
BAB 6 KESIMPULAN DAN SARAN	41
6.1 Kesimpulan	41
6.1.1 Kemampuan Resiko jatuh Pada Lansia	41
6.1.2 Latihan Taichi terhadap Kemampuan Resiko jatuh Pada Lansia.....	41
6.2 Saran.....	42
6.2.1 Bagi Peneliti	42
6.2.2 Bagi Institusi Keperawatan	42
6.2.3 Bagi Masyarakat.....	42
DAFTAR PUSTAKA	43
LAMPIRAN.....	46

DAFTAR TABEL

Tabel 3.1 Kata Kunci pencarian literature latihan taichi untuk kemampuan resiko jatuh pada lansia.....**Error! Bookmark not defined.**

Tabel 3.2 PICOS latihan taichi untuk kemampuan resiko jatuh pada lansia . **Error! Bookmark not defined.**

Tabel 4.1 Hasil Temuan Artikel Latihan Taichi untuk Resiko jatuh pada Lansia**Error! Bookmark not defined.**

Tabel 4.2 Karakteristik Responden Studi.....**Error! Bookmark not defined.**

Tabel 4.3 Resiko jatuh Pada Lanjut Usia**Error! Bookmark not defined.**

Tabel 4.4 Latihan Taichi untuk Kemampuan Resiko jatuh Pada Lanjut Usia**Error! Bookmark not defined.**

DAFTAR GAMBAR

Gambar 2.1 Gerakan Awal.....	15
Gambar 2.2 Gerakan Memeluk Rembulan.....	15
Gambar 2.3 Gerakan Membuka Tirai	16
Gambar 2.4 Gerakan Menyanggah Langit	16
Gambar 2.5 Gerakan Menyanggah Langit dengan Satu Kaki.....	17
Gambar 2.6 Gerakan Tangan Awan.....	17
Gambar 2.7 Gerakan Mengelus Ekor Burung.....	18
Gambar 2.8 Gerakan Memindahkan Gunung Mengeruk Lautan.....	18
Gambar 2.9 Gerakan Pecut Tunggal	19
Gambar 2.10 Gerakan Putri Cantik Menenun.....	19
Gambar 2.11 Gerakan Tendangan Datar Berputar 360 derajat.....	Error! Bookmark not defined.
Gambar 2.12 Gerakan Memutar Roda Kehidupan.....	Error! Bookmark not defined.
Gambar 2.13 Gerakan Tendangan Mengayun	Error! Bookmark not defined.
Gambar 2.14 Gerakan Memungut Batu dan Memetik Daun Muda	Error! Bookmark not defined.
Gambar 2.15 Kerangka Teori Latihan Taichi untuk Resiko jatuh Lansia	Error! Bookmark not defined.
Gambar 3.1 PRISMA Flow Diagram Latihan Taichi Untuk Kemampuan Resiko jatuh Pada Lansia	Error! Bookmark not defined.

BAB 1

PENDAHULUAN

1.1 Latar Belakang

Pada proses kehidupan setiap individu mengalami perkembangan dari sejak lahir hingga lanjut usia (Lansia). Manusia berkembang dari bayi, remaja, dewasa, dan menjadi tua yang merupakan tahap akhir dari proses kehidupan (Permady, 2015). Secara alamiah lansia mengalami berbagai macam kemunduran fungsi tubuh yang menyebabkan lansia mempunyai berbagai masalah (Jayanti, 2016). Saat seseorang memasuki fase lansia, pada saat itu mulai terjadi penurunan fungsi fisiologis yang mengakibatkan terjadinya gangguan degeneratif. Gangguan degeneratif pada lansia salah satunya yaitu penurunan fungsi resiko jatuh dan peningkatan resiko jatuh (Savira, 2016). Jatuh pada lansia merupakan penyebab utama dan sering menyebabkan morbiditas, mortalitas, berkurangnya fungsi, dan menyebabkan cepat masuk ke panti werda pada orang yang berusia 65 tahun ke atas (Kisner, C dan Lynn Allen Colby, 2017). Penyebab kecelakaan yang paling sering terjadi pada orang yang berusia > 65 tahun yaitu jatuh (Abrams dan Berkow, 2013).

Diperkirakan sebanyak 646.000 orang meninggal setiap tahunnya akibat jatuh karena ketidaksengajaan, di mana lebih dari 80% terjadi di negara wilayah Afrika dan Asia Tenggara. Asia Tenggara menyumbang sebanyak 60% kematian. Menurut WHO sebanyak 37,3 juta kejadian jatuh membutuhkan penanganan medis yang intensif setiap tahunnya (WHO, 2018). Di Indonesia proporsi lansia yang mengalami jatuh pada tahun 2019 diperkirakan 1 dari 3 orang berusia 65

tahun ke atas. Faktor penyebab yang paling dominan adalah intrisik (gangguan resiko jatuh) 68% dan faktor lingkungan 31% (Kemenkes RI. 2019). Prevalensi di Indonesia jatuh yang disebabkan cedera yaitu pada usia 65-74 tahun sekitar 67,1% dan pada usia 75 tahun keatas sekitar 78,2% (RISKESDAS, 2018). Kejadian resiko jatuh mengalami peningkatan seiring dengan bertambahnya usia yang lebih dari 65 tahun sebesar 30% dan pada lansia yang berumur lebih dari 80 tahun mencapai 50% (Mupangati, 2018). Prevalensi jatuh di Wisma Sri Tanjung UPT PSTW Banyuwangi diketahui sebesar 26,7% lansia tidak memiliki resiko jatuh, sebesar 53,3% lansia mengalami resiko jatuh rendah dan sebesar 20,0% memiliki resiko jatuh tinggi.

Pada lansia risiko jatuh sering sekali terjadi, dikarenakan pada lansia mengalami penurunan gangguan resiko jatuh, karena adanya fungsi fisiologis yang berubah pada lansia akibat degenerasi dan beberapa merupakan komponen resiko jatuh utama tubuh, seperti visual, ambang rangsang vestibular, kekuatan otot, lingkup gerak sendi, dan sensomotorik. Komponen tersebut merupakan peran penting dalam menjaga kontrol postural pada tubuh. Kontrol postural berfungsi dalam menjaga resiko jatuh tubuh agar tidak mudah terjatuh saat berdiri, berjalan maupun beraktivitas lainnya (D'Silva et al, 2015). Jatuh merupakan keadaan dimana terjadi kegagalan pada manusia dalam mempertahankan resiko jatuh tubuhnya untuk berdiri. Jatuh merupakan masalah fisik yang sering terjadi pada lansia, semakin bertambahnya usia maka terjadi penurunan kondisi fisik, mental dan fungsi tubuh. Risiko jatuh pada lansia merupakan sindrom geriatri yang paling sering terjadi pada lansia (Rudy & Setyanto, 2019). Terdapat dua faktor

yang mempengaruhi risiko jatuh yaitu faktor intrinsik dan faktor ekstrinsik. Faktor intrinsik meliputi kondisi fisik, reflek postural, neuropsikiatrik, perubahan neuromuskular, penurunan visus, gaya berjalan, gangguan jantung dan sirkulasi darah, gangguan sistem anggota gerak, gangguan sistem susunan saraf, dan gangguan psikologis (Rahmawardani, 2018). Faktor ekstrinsik berasal dari lingkungan sekitar lansia yaitu lantai licin dan tidak merata, kursi roda yang tidak terkunci, tersandung oleh benda-benda sekitar, penerangan cahaya yang kurang terang dan gampang terpeleset atau tersandung. Hal tersebut membuat trauma tersendiri bagi lansia sehingga menyebabkan lansia mengalami rasa takut untuk melakukan aktivitas (Rahmawardani, 2018). Gangguan resiko jatuh merupakan penyebab utama yang paling sering mengakibatkan lansia mengalami jatuh (Hwang, 2016). Jatuh dapat mengancam keselamatan jiwa lansia. Jatuh dapat mengakibatkan berbagai jenis cedera, kerusakan fisik dan psikologis. Dampak psikologis adalah syok setelah jatuh dan rasa takut akan terjatuh lagi dapat menimbulkan ansietas yang berlebihan, hilangnya rasa percaya diri, takut, aktivitas sehari-hari menjadi terbatas, fobia atau fobia jatuh (Palvanen et al, 2014). Jatuh juga dapat menimbulkan berbagai komplikasi pada lansia diantaranya sindroma kecemasan setelah jatuh, perlukaan baik jaringan lunak atau patah tulang, disabilitas (Penurunan mobilitas), penurunan status fungsional, penurunan kemandirian, di rawat di Rumah Sakit, peningkatan penggunaan sarana pelayanan kesehatan, dan bahkan bisa sampai meninggal dunia (Murti, Y., 2018).

Resiko jatuh merupakan hal yang sangat penting untuk dimiliki seseorang baik pada usia lanjut maupun pada usia muda agar tetap berada dalam keadaan

yang seimbang dan bisa menyesuaikan diri terhadap gravitasi, permukaan tanah dan ketika melakukan aktivitas sehari-hari agar tidak bergantung pada orang lain (Munawwarah dan Priatna, 2018). Penatalaksanaan yang bisa dilakukan pada lansia dengan risiko jatuh berupa latihan resiko jatuh (Rahmawardani, 2018). Latihan resiko jatuh bertujuan untuk meningkatkan resiko jatuh tubuh, kekuatan otot pada anggota gerak bawah, dan meningkatkan sistem vestibular. Beberapa jenis latihan resiko jatuh yang dapat dilakukan diantaranya senam *Tai Chi*, *Otago Home Exercise Program*, Balance Exercise, Square Stepping Exercise, Ankle Strategy, dan lain-lain (Martins et al, 2018). Senam *Tai Chi* merupakan latihan yang sangat efektif bagi lansia dan di anggap paling aman, praktis, dan hemat dari segi biaya (Mat et al, 2018). Senam *Tai Chi* adalah program latihan resiko jatuh untuk lansia yang didesain khusus untuk mengurangi kejadian jatuh, dengan cara meningkatkan resiko jatuh tubuh, meningkatkan kekuatan otot pada anggota gerak bawah dan memberikan latihan jalan (Mahendra, 2016). Senam *Tai Chi* merupakan program yang hemat dari segi biaya untuk mengurangi jatuh pada orang berusia 65 tahun atau keatas. Program ini dirancang khusus untuk lansia yang terdiri dari 30 menit program latihan resiko jatuh dan latihan penguatan tungkai yang di lakukan di rumah sekitar 3 kali per minggu dan dilengkapi dengan latihan berjalan (Kisner, C dan Allen Colby, 2017). Senam *Tai Chi* menyesuaikan dengan gerakan fungsional sehari-hari pada lansia sehingga juga dapat mengoptimalkan kemampuan lansia dalam melakukan gerakan fungsionalnya.

Resiko jatuh pada lansia dapat ditingkatkan dengan cara latihan fisik yang baik, benar, terukur dan teratur (BBTT) serta latihan yang sesuai dengan tingkat

kesehatan, tingkat aktifitas fisik dan tingkat kebugaran masing-masing individu. Latihan fisik dapat mengurangi resiko kelainan tulang yang menyebabkan resiko jatuh pada lansia. Salah satu latihan fisik yang efektif adalah latihan resiko jatuh taichi. Latihan resiko jatuh taichi ditunjukan untuk membantu meningkatkan kekuatan otot pada anggota gerak bawah (kaki) dan untuk meningkatkan vestibuler atau resiko jatuh tubuh. Berdasarkan hal tersebut latihan resiko jatuh yang dapat dilakukan lansia adalah dengan latihan taichi karena latihan ini sangat membantu untuk mempertahankan tubuhnya agar stabil sehingga mencegah terjatuh yang sering terjadi pada lansia (Nurkoncoro, 2015).

1.2 Rumusan Masalah

Berdasarkan uraian diatas maka rumusan masalah pada penelitian ini bagaimana perbedaan kejadian risiko jatuh sebelum dan sesudah dilakukan terapi senam *Taichi* pada lansia di Wisma Sri Tanjung Panti Sosial Tresna Werdha Banyuwangi.

1.3 Tujuan penelitian

1.3.1 Tujuan Umum

Penelitian ini bertujuan untuk mengetahui perbedaan kejadian risiko jatuh sebelum dan sesudah dilakukan terapi senam *tai chi* pada lansia di Wisma Sri Tanjung Panti Sosial Tresna Werdha Banyuwangi.

1.3.2 Tujuan Khusus

- a) Mengidentifikasi kejadian risiko jatuh lansia sebelum dilakukan terapi *tai chi* pada lansia di Wisma Sri Tanjung Panti Sosial Tresna Werdha Banyuwangi.

- b) Mengidentifikasi kejadian risiko jatuh lansia sesudah dilakukan terapi *tai chi* pada lansia di Wisma Sri Tanjung Panti Sosial Tresna Werdha Banyuwangi.

1.4 Manfaat Penelitian

1.4.1 Bagi Peneliti

Hasil penelitian ini diharapkan dapat dijadikan sebagai bahan masukan atau informasi tambahan tentang pentingnya latihan *tai chi* untuk kejadian risiko jatuh pada lansia.

1.4.2 Bagi Institusi

Hasil penelitian ini dapat digunakan sebagai informasi yang berguna bagi para pembaca untuk menambah wawasan, pengetahuan, dan juga sebagai acuan pembelajaran tentang penerapan asuhan keperawatan terkait dengan latihan *tai chi* untuk kejadian risiko jatuh pada lansia.

1.4.3 Bagi lansia di Panti Sosial Tresna Werdha Banyuwangi

Hasil penelitian ini di harapkan memberikan informasi kepada lansia dengan risiko jatuh lansia bahwa kejadian risiko jatuh dapat diatasi dengan menggunakan pengobatan alternatif yaitu dengan senam *tai chi*.

1.4.4 Bagi UPT Panti Sosial Tresna Werdha Banyuwangi

Dapat memberikan penatalaksanaan pada lansia dengan kejadian risiko jatuh menggunakan terapi senam *tai chi*.

BAB 2

TINJAUAN PUSTAKA

2.1 Konsep Lansia

2.1.1 Definisi Lansia

Lansia atau lanjut usia adalah seseorang yang telah mencapai usia 60 tahun ke atas. Menua bukanlah suatu penyakit, tetapi merupakan proses yang berangsur-angsur mengakibatkan perubahan kumulatif, merupakan proses menurunnya daya tahan tubuh dalam menghadapi rangsangan dari dalam dan luar tubuh. Pada manusia, penuaan berkaitan dengan perubahan degeneratif pada kulit, tulang, pembuluh darah, paru-paru, saraf, dan jaringan tubuh lainnya. Karena kapasitas regenerasi yang terbatas, mereka lebih mungkin menderita untuk menderita berbagai penyakit, sindrom dan penyakit dibandingkan orang dewasa lainnya (Kholifah, 2016).

Lansia secara perlahan akan mengalami penurunan jaringan untuk memperbaiki dan mempertahankan keadaan normal, sehingga lansia sering kali berisiko mengalami berbagai macam penyakit (WHO, 2018). Sebagian besar lansia berusia 60 tahun ke atas akan mulai mengalami perubahan fisik dan mental. Sejak seseorang memasuki usia tua kesehatan fisik akan memburuk dan kualitas hidup menurun seiring dengan berubahnya usia dan status ekonomi (Budi, 2016).

Seseorang yang telah memasuki masa lanjut usia akan terjadi kecenderungan menurunnya berbagai kapasitas fungsional tubuhnya baik pada tingkat selular, maupun pada tingkat organ yang mengakibatkan terjadinya

degenerasi pada proses menua. Hal ini, ini dapat berpengaruh pada perubahan fisiologis secara fisik, fungsi dan persepsi di kehidupan sehari-hari. Setiap individu mengalami perubahan-perubahan pada tubuhnya secara berbeda, ada yang laju penurunannya cepat dan dramatis serta ada juga yang perubahannya lambat. Pada lanjut usia terjadi kemunduran sel-sel karena proses penuaan yang dapat berakibat pada kelemahan organ, kemunduran fisik, timbulnya berbagai macam penyakit (Fauzan; Aldhi, 2019).

2.1.2 Batasan Lanjut Usia

Di Indonesia seseorang dikatakan lanjut usia ketika berumur 60 tahun ke atas. Hal ini sesuai dengan undang-undang nomor 13 tahun 1998 tentang kesejahteraan lanjut usia menekankan hal ini pada Bab 1, pasal 1, ayat 2 (Nugroho, 2008). Usia seorang lansia yang menjadi acuan berbeda hal ini sesuai dengan beberapa pendapat ahli tentang batasan usia lansia adalah sebagai berikut :

1. Menurut organisasi kesehatan dunia (WHO, 2018) ada empat tahapan yaitu:
 - a. Usia pertengahan (middle age) lansia dengan usia 45-59 tahun
 - b. Lansia (elderly) lansia dengan usia 60-74 tahun
 - c. Lansia tua (old) lansia dengan usia 75-90 tahun
 - d. Usia sangat tua (very old) lansia dengan usia >90 tahun
2. Menurut (Kemenkes RI, 2019) pembagian usia lanjut ada 3 yaitu :
 - a. Kelompok lansia dini (45 sampai dengan <60 tahun) merupakan kelompok yang baru memasuki lansia atau pra lansia
 - b. Kelompok lansia usia pertengahan (60-70 tahun)

- c. Kelompok lansia yang berisiko tinggi yaitu lansia yang berusia >70 tahun
3. Menurut (Elipoulus, 2010) batasan usia lanjut dibagi menjadi 3 yaitu :
- a. Setengah tua yaitu lansia berusia antara 60-74 tahun
 - b. Tua yaitu lansia yang berusia antara 75-100 tahun
 - c. Sangat tua yaitu lansia yang berusia >100 tahun

2.2 Konsep Resiko jatuh Tubuh

Menurut (Suhartono, 2006) bahwa resiko jatuh tubuh adalah kemampuan seseorang untuk mempertahankan posisi tubuh agar tetap statik dan dinamik. Dapat disimpulkan yang dimaksud dengan resiko jatuh tubuh adalah kemampuan tubuh mempertahankan postur agar tetap tegak dengan gravitasi dan tetap dalam landasan penopang untuk mengatur seluruh aktivitas diam (static) atau sedang bergerak (dinamik).

2.2.1 Perubahan Resiko jatuh Tubuh Dalam Proses Menua

Proses menua adalah perubahan yang berkaitan dengan berjalanya waktu dan bersifat universal, intrinsik dan progresif. Perubahan tersebut mengakibatkan terjadinya penurunan kemampuan sel dan jaringan beradaptasi dengan lingkungan dan bertahan hidup. Perubahan pada fungsi fisiologis yaitu terjadi perubahan pada sensori, neurologis dan sistem saraf pusat, motorik dan muskuluskeletal (Achamanegara, 2012)

Proses penuaan berkaitan dengan penurunan sistem organ salah satunya sistem muskuluskeletal, menurunnya sistem muskuluskeletal berdampak pada sistem resiko jatuh tubuh karena terjadi atrofi otot yang menyebabkan penurunan kekuatan otot, terutama pada otot ekstremitas bawah sehingga terlambat mengantisipasi bila terjatuh atau terpeleset. Dengan terganggunya sistem muskuluskeletal akan berdampak pada strategi postural tubuh sehingga mempengaruhi resiko jatuh (Maryam, 2010)

2.2.2 Komponen Resiko jatuh Tubuh

Penuaan ditandai dengan perubahan komponen dari sistem muskuluskeletal yang berpengaruh terhadap resiko jatuh. Konsekuensi dari penuaan adalah pada penurunan resiko jatuh yang dimanifestasikan dalam beberapa komponen yaitu berdiri, bersandar, dan membungkuk, performa gerakan terkontrol dan respon terhadap gangguan eksternal (Achamanegara, 2012).

2.2.3 Penilaian Fungsi Resiko jatuh

Ada berbagai jenis instrumen yang digunakan untuk melakukan penilaian terhadap resiko jatuh. Salah satu instrumen yang memiliki tingkat validitas dan reliabilitas yang tinggi adalah BBS. BBS adalah sebuah skala yang terdiri dari 14 item observasi yang digunakan untuk menilai resiko jatuh lansia di komunitas (Bitensky, 2008).

2.2.4 Faktor-Faktor yang Mempengaruhi Resiko jatuh Tubuh Lansia

a. Usia Lansia

Dibandingkan dengan orang yang lebih muda, menunjukkan tingkat ketidakseimbangan yang lebih besar ketika berdiri yang ditandai dengan

goyangan postur tubuh. Kebanyakan penelitian menunjukkan bahwa ketidakseimbangan meningkat dengan bertambahnya usia semakin bertambah usia sistem tersebut semakin menurun (Achmanegara, 2012).

b. Jenis Kelamin

(Tedeiksar, 2017)mengatakan lansia perempuan menunjukkan angka ketidakseimbangan lebih besar daripada laki-laki. sehingga pada lansia perempuan cenderung akan mengalami risiko jatuh. Hasil penelitian maryam 17 diperoleh bahwa pada lansia perempuan kurang dalam melakukan aktivitas fisik dibandingkan dengan lansia laki-laki sehingga dapat mempengaruhi resiko jatuh tubuh.

c. Aktivitas Fisik

Aktivitas fisik dapat mempertahankan fungsi muskuluskeletal sehingga resiko jatuh tubuh pada lansia dapat dipertahankan. (Maryam, 2009) menyebutkan bahwa pada lansia yang memiliki aktivitas yang kurang berisiko terjadi gangguan resiko jatuh dari pada lansia yang aktivitasnya baik. Aktivitas yang teratur dapat meningkatkan kebugaran, kekuatan dan koordinasi serta resiko jatuh tubuh pada lansia (Harsuki, 2003). Aktivitas fisik mempunyai dampak positif terhadap resiko jatuh tubuh, serta menurunkan resiko jatuh (SKelton, 2011)

d. Obat-obatan dan Alkohol

Obat-obatan tertentu mempengaruhi gangguan resiko jatuh tubuh pada lansia karena menimbulkan efek mengantuk dan lansia menjadi kurang waspada. Beberapa diantaranya obat sedatif. Lansia dengan konsumsi obat

yang banyak dapat mempengaruhi resiko jatuh. Alkohol dapat menurunkan kewaspadaan dan dapat mempengaruhi resiko jatuh tubuh lansia(Savira, 2016).

2.3 Konsep Senam *Tai Chi*

2.3.1 Pengertian Senam *Tai Chi*

Senam merupakan latihan tubuh yang dapat membentuk dan mengembangkan pribadi secara harmonis dengan terdapat beberapa unsur seperti melompat, memanjat, dan resiko jatuh (Widiantri & Proverawati , 2012). Taichi merupakan suatu bentuk gaya seni bela diri dari china yang terdiri atas latihan meditasi, pergerakan melingkar, peregangan yang halus dan posisi seimbang pada tubuh. Taichi merupakan suatu keterampilan, kemampuan, atau usaha yang pada umumnya diterapkan pada pelatihan bela diri (Serikali, 2006). Senam taichi pada prinsipnya berbeda dengan senam yang lainnya seperti membakar kalori, membuat jantung berdebar-debar atau mengecilkan perut yang membedakan adalah senam ini melatih kekuatan otot (Sutanto, 2013). Senam taichi dapat dilakukan sebanyak 3 kali dalam seminggu selama sebulan dengan hasil yang efektif (Putri, 2014). Senam taichi juga dapat dengan efektif dilakukan selama 3 minggu dengan frekuensi 3 kali dalam seminggu (Ismiati, 2013).

2.3.2 Tujuan Senam *Tai Chi*

Tujuan dari senam taichi adalah untuk mengembangkan hubungan yang lebih erat antara tubuh, pikiran dan jiwa bagi pelaksana senam taichi ini. Senam taichi ini dilakukan dengan teknik gerakan yang sangat lambat untuk menyebarkan energi dalam tubuh. Taichi sangat berpengaruh pada tubuh dalam

hal resiko jatuh dan ketenangan. Tujuan senam taichi menurut (Sutanto, 2013) yaitu sebagai berikut:

- a. Dapat diaplikasikan kepada anak-anak hingga manula untuk meningkatkan kesehatan fisik, resiko jatuh jiwa dan mental serta memperkuat daya tahan tubuh yang merupakan suatu akibat dari adanya meditasi dalam senam sehingga mengaktifkan sistem endokrin serta pelepasan neurotransmitter.
- b. Mengurangi kecemasan dan depresi
- c. Memperbaiki resiko jatuh, fleksibilitas dan kekuatan otot
- d. Mengurangi resiko jatuh
- e. Memperbaiki kualitas tidur
- f. Menstabilkan tekanan darah sebab terdapat unsur meditasi, teknik relaksasi yang berdampak baik bagi denyut jantung
- g. Memperbaiki kapasitas jantung pada lansia
- h. Menghilangkan nyeri kronik pada gangguan neuromuskuluskeletal
- i. Meningkatkan kapasitas energi
- j. Meningkatkan kemampuan antioksidan dan imunitas
- k. Mencegah osteoporosis
- l. Merangsang organ internal untuk dapat berjalan dengan baik dan
- m. Mempertahankan kualitas hidup yang maksimal

2.3.3 Prinsip Senam *Tai Chi*

Terdapat 10 prinsip dalam melakukan gerakan *Tai Chi* yaitu sebagai berikut (Sutanto, 2013):

- a. Titik berat atau gravitasi tubuh terletak pada beberapa centimeter dibawah pusat (Tan Tiem);
- b. Gerakan yang diberikan terus sambung menyambung tanpa putus;
- c. Menggunakan pernafasan perut bukan pernafasan dada
- d. Pikiran masuk ke dalam titik berat atau gravitasi tubuh
- e. Terdapatnya gerakan membuka (ekspansi) dan menutup (kontraksi), maju mundur, naik turun, clockwise serta counter clockwise dan nafas yang masuk serta keluar secara berkesinambungan.
- f. Gerakannya yang melingkar seperti ulir
- g. Melatih rasa bukan otot;
- h. Merupakan satu kesatuan yang dimana satu anggota tubuh bergerak yang lain juga ikut bergerak;
- i. Pada saat bergerak mencari suatu ketenangan dan begitu juga sebaliknya; serta
- j. Gerakannya yang lincah.

2.3.4 Penatalaksanaan Senam Tai Chi

Gerakan-gerakan yang terdapat di dalam Senam *Tai Chi* ini adalah sebagai berikut (Sutanto, 2010)

No	Gambar Gerakan	Keterangan
1	 Gambar 2.1 Gerakan Awal	Posisi awal, yaitu tubuh dalam posisi tegak lalu lutut menekuk bersamaan dengan kedua tangan yang diangkat membentuk sudut 45 derajat mendekati dada. Gerakan dilakukan secara perlahan lalu diturunkan secara bersamaan;
2	 Gambar 2.2 Gerakan kedua	Posisi kedua tangan di angkat ke atas secara bersamaan lalu diturunkan secara bersamaan dengan kedua kaki di tekukuk. Lakukan secara perlahan lalu kembali ke posisi semula

3	 Gambar 2.3 Gerakan ketiga	Membuka tirai kehidupan, yaitu kedua tangan mengumpul diperut naik hingga ke dada lalu melakukan gerakan menyamping hingga tangan sejajar dengan lengan. Lakukan gerakan dengan perlahan lalu kembali ke posisi semula;
4	 Gambar 2.4 Gerakan kelima	Menarik kedua bahu kebelakang secara bersamaan dengan posisi tangan di tekuk di depan dan kedua kaki ditekuk. sebaliknya. Kembali ke posisi awal;

5	 Gambar 2.5 Gerakan kelima	Menarik kedua bahu kedepan secara bersamaan dengan posisi tangan di tekuk di depan dan kedua kaki ditekuk. sebaliknya. Kembali ke posisi awal;
6	 Gambar 2.6 Gerakan ke enam	Tangan awan, yaitu posisi tubuh menghadap ke kanan dengan tangan kanan sejajar dengan dada membentuk sudut 45 derajat sedangkan tangan kiri berada di pusat menghadap ke atas. Kaki membuka dengan kaki kanan menghadap ke kanan dan kaki kiri menghadap ke depan. Tangan kanan bergeser ke arah kiri begitu juga dengan kaki bergeser mengikuti gerak tubuh menghadap ke kiri. Lakukan sebaliknya dengan membelokkan tubuh ke arah yang berlawanan.
7		Mengelus ekor burung, tubuh dalam posisi menghadap ke arah kanan dengan kaki

	 Gambar 2.7 Gerakan ketujuh	menekuk lalu kaki kanan maju ke depan menjadi tumpuan tubuh sedangkan tangan kiri ke belakang dan tangan kanan ke arah ke depan dan atas. Lalu tarik kedua tangan hingga berkumpul, dan lakukan kepada kaki kiri dengan gerakan yang sama. Lalu kembali ke posisi semula
8	 Gambar 2.8 Gerakan ke delapan	Gerakkan kedua tangan ke samping kanan dengan posisi kaki kanan maju ke menyamping sebagai tumpuan tubuh. lakukan pada kaki kiri dengan gerakan tangan yang. Lakukan pada arah yang sebaliknya pula

9	 Gambar 2.9 Gerakan kesembilan	Gerakkan tangan kanan ke arah samping kiri dengan posisi kaki kanandi tekuk sebagai tumpuan. Lakukan pada tangan kiri ke kanan dengan gerakan yang sama . kembali pada posisi semula
10	 Gambar 2.10 Gerakan penutup	Gerakan kedua tangan di angkat ke atas dari samping sampai bertemu kedua tangan di dada dengan posisi kedua kaki tegak lurus dan rapat.

No	Gambar Gerakan	Keterangan
1	 Gambar 2.1 Gerakan Awal	Posisi awal, yaitu tubuh dalam posisi tegak lalu lutut menekuk bersamaan dengan kedua tangan yang diangkat membentuk sudut 45 derajat mendekati dada. Gerakan dilakukan secara perlahan lalu diturunkan secara bersamaan;

2	 Gambar 2.2 Gerakan kedua	Posisi kedua tangan di angkat ke atas secara bersamaan lalu diturunkan secara bersamaan dengan kedua kaki di tekukuk. Lakukan secara perlahan lalu kembali ke posisi semula
3	 Gambar 2.3 Gerakan ketiga	Membuka tirai kehidupan, yaitu kedua tangan mengumpul diperut naik hingga ke dada lalu melakukan gerakan menyamping hingga tangan sejajar dengan lengan. Lakukan gerakan dengan perlahan lalu kembali ke posisi semula;

BAB 3

TINJAUAN KASUS

3.1 Pengkajian

3.1.1 Data Inti

a) Sejarah

Pada tahun 1961 pemerintah Indonesia menyerahkan aset Negara tersebut kepada Dinas Sosial Provinsi Jawa Timur sebagai Panti Multi kegiatan permasalahan sosial dilakukan sampai dengan tahun 1964 dan kurun waktu tersebut yang menjadi Pimpinan bernama Bapak ISHAJI. Pada tahun 1965 menjadi Panti Aneka Permasalahan Sosial yang menangani bermacam-macam permasalahan sosial di wilayah kabupaten Banyuwangi khususnya dan wilayah timur pada umumnya. Kegiatan tersebut berlangsung sampai dengan tahun 1969.


Pada tahun 1970 – 2002 berubah fungsi menjadi Panti Karya WISMA BHAKTI lokasi jauh sebagai Unit Pelayanan Sosial yang spesifikasi pelayanannya adalah penanganan gelandangan dan pengemis. Pada tahun 2002 – 2004 berubah menjadi UPS Bina Karya dibawah UPT Rehsos Gepeng Pasuruan. Pada tahun 2009 berubah beralih fungsi menjadi Unit Pelayanan Teknis (UPT) Pelayanan Sosial Tresna Werdha (PSTW) Banyuwangi dengan spesifikasi pelayanannya adalah penanganan Lanjut Usia terlantar.

b) Demografi

1) Komposisi Penduduk

Berdasarkan jumlah dari 100 lansia di PSTW Banyuwangi, lansia yang berada di Wisma Sri Tanjung berjumlah 15 lansia, berdasarkan jenis kelamin adalah sebagai berikut:


No	Jenis Kelamin	Jumlah
1	Laki-laki	19
2	Perempuan	0
Total jumlah lansia		19



Berdasarkan diagram diatas didapatkan hasil presentase jenis kelamin perempuan 100%

2) Jumlah Lansia Berdasarkan Usia Lansia Menurut WHO


No.	Usia	Jumlah
1	45-59	-
2	60-74	8
3	75-90	7
4	>90	-
Jumlah		15



Berdasarkan diagram diatas didapatkan hasil presentase usia 75-90 sebesar 53% dan usia 60-74 sebesar 42%.

3) Status Perkawinan


No	Status Perkawinan	Jumlah
1	Kawin	-
2	Tidak Kawin	-
3	Janda / Duda	15
Jumlah		15



Berdasarkan diagram diatas didapatkan hasil presentase status perkawinan dengan janda/duda 68%, kawin 21%, dan tidak kawin 11%.

4) Agama


No	Agama	Jumlah
1	Islam	17
2	Kristen	2
3	Hindu	-
4	Budha	-
5	Konghucu	-
Jumlah		19



Berdasarkan diagram diatas didapatkan hasil presentase agama yang dianut di wisma Minak Jinggo Putra sebanyak 89% beragama islam dan 11% beragama kristen

5) Status Kesehatan

No.	Penyakit	Jumlah
1	ISPA	2
2	TBC	-
3	Hipertensi	9
4	Jantung	-
5	Ginjal	-
6	Stoke	1
7	Diabetes Milletus	-
8	DHF	-
9	Diare	1
10	Gatal-gatal	5
11	Gangguan Jiwa	1
12	Lain-lain	-
Jumlah		19



Berdasarkan diagram diatas didapatkan hasil presentase status kesehatan lansia di wisma Minak Jinggo Putra penyakit yang diderita lansia yaitu hipertensi 48%, gatal-gatal 26%, ISPA 11%, stroke 5, diare 5%.

3.1.2 Data Subsystem

a) Lingkungan Fisik

1) Wisma Sri Tanjung

Wisma Sri Tanjung adalah wisma yang terletak di sebelah kiri belakang PSTW Banyuwangi, Wisma Sri Tanjung merupakan tempat tinggal lansia perempuan yang mandiri. Wisma Sri Tanjung memiliki 16 bed/tempat tidur sedangkan yang terisi sejumlah 15 bed/tempat tidur. Pencahayaan di Wisma Sri Tanjung cukup. Jenis bangunan wisma yaitu permanen. Jenis lantai wisma yaitu keramik. Ventilasi udara lebih dari 10% dari luas bangunan. Jumlah ruangan di Wisma Sri Tanjung ruangan luasnya $\pm 3 \times 6 \text{ m}^2$

dan ruangan. Sumber air minum yaitu air galon isi ulang. Terdapat 5 kamar mandi, jenis jamban yaitu menggunakan WC leher angsa.

2) Perilaku Sehat


i. Kebiasaan Mandi dan Ganti Pakaian

No	Keterangan	Jumlah
1	Rutin	15
2	Tidak Rutin	-
Jumlah Lansia		15

Keterangan:

Rutin: mandi 2 x sehari dan berganti pakaian


Tidak Rutin: dalam 1 hari kadang mandi, kadang tidak mandi sama sekali dan tidak berganti pakaian



Berdasarkan diagram diatas didapatkan hasil presentase kebiasaan mandi dan ganti pakaian di wisma Minak Jinggo Putra sebanyak 67% tidak rutin dan sebanyak 33% rutin

ii. Kebiasaan Merokok

No.	Keterangan	Jumlah
1.	Merokok	1
2.	Tidak Merokok	14
Jumlah Lansia		15



Berdasarkan diagram diatas didapatkan hasil presentase kebiasaan merokok di wisma Minak Jinggo Putra sebanyak 79% merokok dan 21% tidak merokok

iii. Kebiasaan Makan Sayur dan Lauk Pauk

No.	Keterangan	Jumlah
1.	Ya	15
2.	Tidak	0
Jumlah Lansia		15



Berdasarkan diagram diatas didapatkan hasil presentase kebiasaan lansia mengkonsumsi sayur dan lauk pauk sebesar 100%.

iv. Kebiasaan Olahraga


No.	Keterangan	Jumlah
1.	Rutin Olahraga	6
2.	Tidak Rutin Olahraga	9
Jumlah Lansia		15

Keterangan:

Rutin Olahraga: mengikuti senam yang di adakan oleh UPT

PSTW Banyuwangi tiap hari selasa dan jum'at.

Tidak Rutin Olahraga: tidak megikuti senam yang di adakan oleh UPT PSTW Banyuwangi tiap hari selasa dan jum'at.



Berdasarkan diagram diatas didapatkan hasil presentase kebiasaan olahraga lansia di wisma Minak Jinggo Putra yang rutin mengikuti olahraga sebanyak 74% dan yang tidak rutin mengikuti olahraga sebanyak 26%.


3) Sumber Air

- a. Sumber air minum dan masak di UPT PSTW Banyuwangi menggunakan Air Galon isi ulang
- b. Sumber air mandi dan cuci baju di UPT PSTW Banyuwangi menggunakan air sumber/sumur yang ditampung di tandon.

4) Pembuangan Sampah

Pembuangan sampah di UPT PSTW Banyuwangi diantaranya :

No.	Keterangan	Jumlah
1	Ditimbun	-
2	Dibakar	3
3	TPA	2
Jumlah		5



5) Pembuangan Limbah

Pembuangan limbah di UPT PSTW Banyuwangi diantaranya:

No.	Keterangan	Jumlah
1.	Sungai	-
2.	Got	1
3.	Tidak Ada	-
Jumlah		1


b) Keamanan

No	Fasilitas Keamanan	Jumlah
1	Pemadam Kebakaran	-
2	Pos Polisi	-
3	Pos Satpam	1
	Lain-lain	-

Sistem keamanan yang ada di UPT PSTW Banyuwangi terbilang kondusif dimana kesadaran lansia untuk ikut menjaga keamanan PSTW terbilang cukup baik adapun sarana prasarana yang ada di UPT PSTW Banyuwangi adanya pos satpam yang terletak di depan pintu masuk PSTW Banyuwangi yang menjadi central keamanan di UPT PSTW Banyuwangi.

c) Pelayanan Kesehatan

No.	Keterangan	Jumlah
1.	Rutin	15
2.	Tidak Rutin	-
3.	Tidak Pernah	-
Jumlah		15



Berdasarkan diagram diatas didapatkan hasil presentase pemeriksaan rutin yang dilakukan lansia di wisma Minak Jinggo Putra sebanyak 85% lansia rutin memeriksa kesehatan dan sebanyak 15% lansia tidak rutin memeriksa kesehatannya.

d) Ekonomi

Sumber pendapatan ekonomi yang diperoleh berasal dari Dinas Pemerintah Provinsi Jawa Timur.

e) Politik dan Pemerintahan

UPT PSTW Banyuwangi dinaungi langsung oleh pemerintah Provinsi Jawa Timur.

f) Sistem Komunikasi

Sistem komunikasi antar wisma berjalan dengan baik, tiap-tiap wisma memiliki ketua untuk menyampaikan keluhan kesah para lansia, dan para ketua langsung menyampaikan pada petugas UPT PSTW Banyuwangi.

g) Pendidikan

Pendidikan rata rata di Wisma Sri Tanjung adalah tamat SMA.

h) Rekreasi

UPT PSTW Banyuwangi terdapat sarana aktivitas rekreasi untuk lansia diantaranya:

No.	Fasilitas Rekreasi	Jumlah
1	TV	4
2	Radio	0
3	Taman (gardu)	3

3.1.3 Persepsi

a) Persepsi petugas

Setiap petugas memiliki pengaruh paling besar terhadap apa yang dikerjakan dan kebutuhan lansia yang harus dipenuhi. Dapat

disimpulkan bahwa petugas harus lebih memahami tentang peran dan fungsinya sebagai pemberi pelayanan pada lansia. Direkomendasikan agar yang berwenang dapat menempatkan petugas UPT PSTW Banyuwangi sesuai dengan peran dan fungsi serta kewenangannya.

b) Persepsi lansia

Setiap lansia satu sama lain mempunyai persepsi yang tidak sama tentang keberadaan lingkungan fisik dan sosial di UPT PSTW Banyuwangi. Latar belakang yang dimiliki lansia tentulah berbeda, baik latar belakang keluarga, lingkungan, tempat tinggal, status sosial ekonomi, dan karakter lansia itu sendiri. Dari latar belakang yang berbeda-beda, akan memberikan implikasi yang tidak sama terhadap persepsi atau pandangan lansia tentang UPT PSTW Banyuwangi.

BAB 4

3.2 Hasil

Berdasarkan hasil pengkajian sebelum dilakukan intervensi pada lansia di Wisma Sri Tanjung diketahui bahwa sebagian besar lansia memiliki resiko jatuh, lansia yang memiliki resiko jatuh merupakan lansia dengan masalah fisiologis pada ekstermitas dan memiliki masalah kesehatan seperti osteoarthritis, gout arthritis dan banyak masalah kesehatan lain yang menyebabkan penurunan fungsi fisiologis pada ekstermitas. Saat dilakukan pengkajian diketahui bahwa lansia memiliki riwayat jatuh 3 bulan terakhir cukup tinggi, lansia yang berisiko jatuh cukup tinggi memiliki lebih dari satu masalah kesehatan. Setelah dilakukan pengkajian lansia memiliki kriteria risiko jatuh diantaranya tidak beresiko jatuh 4, beresiko jatuh rendah 8, dan resiko jatuh tinggi 3 Lansia di Wisma Sri Tanjung sebagian menggunakan alat bantu jalan berupa kruk atau tongkat. Berdasarkan hasil pengamatan diketahui bahwa gaya berjalan lansia sebagian besar lemah dan lansia menyadari masalah fisik yang dialami.

Penelitian ini dilakukan dengan data primer lansia yang berada di wisma Sri Tanjung UPT PSTW Banyuwangi. Jumlah subjek pada penelitian ini adalah 15 orang. Dari hasil pengumpulan data didapatkan jenis kelamin, usia, efektifitas senam taichi sebelum dan sesudah dilakukan pada lansia dengan resiko jatuh.

Paired Samples Test

		Paired Differences			
		Mean	Std. Deviation	Std. Error Mean	95% Confidence ...
					Lower
Pair 1	sebelum diberikan perlakuan - setelah dibeikan perlakuan	,00000	1,73205	1,00000	-4,30265

Paired Samples Test

		Paired ...	t	df	Sig. (2-tailed)
		95% Confidence ...			
		Upper			
Pair 1	sebelum diberikan perlakuan - setelah dibeikan perlakuan	4,30265	,000	2	1,000

1. Hasil intervensi latihan Taichi terhadap resiko jatuh pada lansia setelah 3 kali pertemuan selama seminggu didapatkan hasil bahwa latihan taichi memiliki pengaruh yang efektif terhadap resiko jatuh yang dialami lansia di Wisma Sri Tanjung. Setelah intervensi didapatkan skor jatuh yang diukur menggunakan *Fall Morse Scale* cukup menurun menjadi tidak beresiko jatuh 6, beresiko jatuh rendah 7 dan resiko jatuh tinggi 2.

BAB V

PEMBAHASAN

5.1 Risiko Jatuh Lansia Sebelum Dilakukan terapi Tai Chi

Berdasarkan hasil pengkajian sebelum dilakukan intervensi pada lansia di Wisma Sri Tanjung diketahui bahwa sebagian besar lansia memiliki resiko jatuh, lansia yang memiliki resiko jatuh merupakan lansia dengan masalah fisiologis pada ekstermitas dan memiliki masalah kesehatan seperti osteoarthritis, gout arthritis dan banyak masalah kesehatan lain yang menyebabkan penurunan fungsi fisiologis pada ekstermitas. Saat dilakukan pengkajian diketahui bahwa lansia memiliki riwayat jatuh 3 bulan terakhir cukup tinggi, lansia yang berisiko jatuh cukup tinggi memiliki lebih dari satu masalah kesehatan. Lansia di Wisma Sri Tanjung sebagian menggunakan alat bantu jalan berupa kruk atau tongkat. Berdasarkan hasil pengamatan diketahui bahwa gaya berjalan lansia sebagian besar lemah dan lansia menyadari masalah fisik yang dialami.

Penurunan yang terjadi pada fisiologis tubuh makin bertambah seiring bertambahnya usia, terutama yang berpengaruh pada pengontrol resiko jatuh. Resiko jatuh yang kurang baik pada lansia dapat mengakibatkan jatuh. Hal ini sejalan dengan teori yang menyatakan bahwa keseimbangan merupakan kontrol motor yang kompleks melibatkan deteksi dan integrasi informasi sensori untuk memeriksa posisi dan gerakan tubuh dalam ruang dan eksekusi respons muskulokeletal yang

tepat untuk mengontrol posisi tubuh dengan konteks lingkungan dan tugas. Oleh karena itu, kontrol resiko jatuh memerlukan interaksi sistem nervous dan muskuloskeletal dan pengaruh kontekstual (Widarti, 2020). Pada penelitian yang dilakukan di Taiwan diketahui bahwa resiko jatuh pada lansia mengalami penurunan seiring dengan berjalan nya waktu. Penelitian yang dilakukan juga menyimpulkan bahwa beberapa latihan fisik dapat membantu lansia menjaga resiko jatuh mereka (Wang, et al., 2016).

Keseimbangan memiliki peran yang penting bagi lansia, karena keseimbangan merupakan sebuah kontrol dari system gerak lansia yang kompleks hal tersebut melibatkan integrasi informasi sensori dan respon yang baik. Oleh karena itu beberapa peneliti melakukan penelitian tentang resiko jatuh dengan cara melakukan berbagai rangkaian tes seperti kemampuan resiko jatuh lansia dalam turun tangga, kemudian dilakukan tes berdiri dengan satu kaki hal tersebut juga dilakukan untuk mengetahui resiko jatuh yang ada pada lansia.

5.2 Latihan Taichi terhadap Resiko jatuh pada Lanjut Usia

Hasil intervensi latihan Taichi terhadap resiko jatuh pada lansia setelah 3 kali pertemuan selama seminggu didapatkan hasil bahwa latihan taichi memiliki pengaruh yang efektif terhadap resiko jatuh yang dialami lansia di Wisma Sri Tanjung. Setelah intervensi didapatkan skor jatuh yang diukur menggunakan *Fall Morse Scale* cukup menurun. Latihan taichi yang dilakukan lansia

Masalah resiko jatuh sering kali mengancam kualitas hidup lansia, karena lansia yang mengalami penurunan resiko jatuh akan memiliki ketergantungan fisik pada orang lain. Jatuh merupakan salah satu penyebab multifaktorial, faktor resiko jatuh, penurunan resiko jatuh serta penurunan kognitif terjadi karena terkait usia (Kim, 2019). Pada lansia kehilangan resiko jatuh sering kali terjadi diakibatkan oleh sarcopenia dan kerapuhan (Landi, et al., 2012). Untuk menangani masalah resiko jatuh dikenal berbagai macam latihan yang dapat dilakukan lansia salah satunya menggunakan latihan taichi. Taichi dilakukan dengan gerakan lembut, tidak memaksa dan terkoordinasi adalah bentuk latihan yang tepat dilakukan oleh lansia karena gerakan dilakukan tanpa perlu memaksa gerak persendian dan system kardiovaskular (Huang & Liu, 2015). Senam tai chi berasal dari seni bela diri Cina kuno. Aktivitas fisik ini mempraktikkan rangkaian gerakan yang mesti dilakukan secara perlahan dan penuh konsentrasi, disertai dengan teknik bernapas dalam. Dalam pelaksanaan latihan taichi perlu diperhatikan terkait waktu dan durasi pemberian latihan taichi terutama pada lansia, idealnya waktu dilakukannya latihan hanya selama kurang lebih 30 menit (Huang & Liu, 2015). Disebut juga tai chi chuan, olahraga ini bersifat nonkompetitif dan lebih menitikberatkan pada latihan fisik individual serta peregangan. Sehingga latihan taichi sangat sesuai dengan kebutuhan aktivitas fisik yang dianjurkan bagi lansia karena setiap gerakan dalam senam tai chi akan berlanjut ke gerakan berikutnya dalam aliran yang lembut tanpa jeda,

untuk memastikan tubuh tidak berhenti bergerak (Wang, et al., 2016). Senam tai chi terdiri dari beberapa aliran. Tiap aliran memiliki prinsip gerakan dan metode latihan masing-masing. Ada aliran tai chi yang berfokus pada menjaga kebugaran dan kesehatan, dan ada juga aliran yang lebih berfokus pada sisi seni bela diri (Day, et al., 2015). Berdasarkan penelitian yang dilakukan tentang taichi pada lansia. Diketahui bahwa taichi yang dilakukan secara rutin dapat mengatasi berbagai macam masalah kesehatan yang terjadi pada lansia seperti masalah pada sendi, pada ekstermitas dan anggota gerak tubuh lainnya. Selain itu latihan taichi juga berpengaruh kepada fungsi kognitif seseorang terutama lansia, karena secara fisiologis lansia sudah mengalami penurunan kognitif tetapi dengan taichi terbukti mampu mengurangi efek penurunan kognitif lansia (Miller & Taylor-Piliae, 2014). Pada penelitian yang pernah dilakukan menunjukkan bahwa taichi dapat meningkatkan kekuatan otot di ekstermitas bawah dan meningkatkan kontrol resiko jatuh, proprioception dan adaptasi postural sehingga mengurangi resiko jatuh dan meningkatkan resiko jatuh (Huang, et al., 2017).

Latihan taichi memiliki gerakan yang sangat sederhana dan tidak memerlukan banyak gerakan juga dinilai sangat sesuai dengan kemampuan lansia dalam melakukan aktivitas fisik sehingga latihan taichi dinilai sesuai dengan kebutuhan lansia. Latihan taichi secara teratur setidaknya setiap sesi 30 menit dan dilakukan selama kurang lebih 3 kali seminggu memiliki efek yang baik pada kesehatan lansia, taichi akan

meningkatkan resiko jatuh lansia selain itu taichi juga bermanfaat mengurangi resiko jatuh dan masalah kognitif pada lansia.

BAB VI

KESIMPULAN DAN SARAN

6.1 Kesimpulan

Berdasarkan analisis dari beberapa artikel tentang latihan taichi untuk resiko jatuh lansia dapat disimpulkan:

6.1.1 Resiko Jatuh Pada Lansia Sebelum Latihan Taichi

Sebelum dilakukan intervensi pada lansia di Wisma Sri Tanjung diketahui bahwa sebagian besar lansia memiliki resiko jatuh, lansia yang memiliki resiko jatuh merupakan lansia dengan masalah fisiologis pada ekstermitas dan memiliki masalah kesehatan Taichi yang dilakukan secara rutin dapat mengatasi berbagai macam masalah kesehatan yang terjadi pada lansia seperti masalah pada sendi, pada ekstermitas dan anggota gerak tubuh lainnya. Selain itu latihan taichi juga berpengaruh kepada fungsi kognitif seseorang terutama lansia.

6.1.2 Latihan Taichi terhadap Resiko jatuh pada Lanjut Usia

Latihan taichi secara teratur setidaknya setiap sesi 30 menit dan dilakukan selama kurang lebih 3 kali seminggu memiliki efek yang baik pada kesehatan lansia, taichi akan meningkatkan resiko jatuh lansia selain itu taichi juga bermanfaat mengurangi resiko jatuh dan masalah kognitif pada lansia. Bentuk latihan taichi yang sangat sederhana dan tidak memerlukan banyak gerakan juga dinilai sangat sesuai dengan kemampuan lansia dalam melakukan aktivitas fisik sehingga latihan taichi dinilai sesuai dengan kebutuhan lansia.

6.2 Saran

Berdasarkan hasil penelitian diatas maka peneliti menyarankan beberapa hal sebagai berikut:

6.2.1 Bagi Peneliti

Diharapkan *mini research* ini dapat menambah pengetahuan terkait latihan taichi dan resiko jatuh pada lansia serta dapat diaplikasikan pada saat melakukan asuhan keperawatan pada lansia.

6.2.2 Bagi Institusi

Diharapkan hasil *mini research* ini dapat menambah bahan refrensi bagi institusi pendidikan mengenai terapi latihan taichi sebagai salah satu terapi alternatif untuk meningkatkan resiko jatuh pada lansia.

6.2.3 Bagi Masyarakat

Diharapkan masyarakat yang mengalami masalah resiko jatuh dapat menerapkan latihan taichi sebagai salah satu terapi alternatif untuk menurunkan masalah resiko jatuh yang dapat berakibat resiko jatuh pada lansia.

DAFTAR PUSTAKA

- Achamanegara. (2012). *Faktor Mempengaruhi Resiko jatuh Tubuh*. Jakarta: Jakarta Gemilang.
- Budi, R. S. (2016). *gambaran aktivitas fisik dengan kualitas tidur lansia*. Yogyakarta: UNY.
- Chen, P.-J., Penn, I.-W., Wei, S.-H., Chuang, L.-R., Sung, W.-H., & Sung. (2020, May 20). Augmented reality-assisted training with selected Tai-Chi movements improves balance control and increases lower limb muscle strength in older adults: A prospective randomized trial. *Journal of Exercise Science & Fitness*, 142-147. doi:https://doi.org/10.1016/j.jesf.2020.05.003
- Chen, S. (2019). Effect of Tai Chi Exercises on the Balance, Functional Gait, and Flexibility of Elderly Filipino Males. *Advances in Social Science, Education and Humanities Research*, 268, 119.
- Day, I., Hill, K., Stathakis, VZ, Hoke, M., & Piliae. (2015). Impact of tai-chi on falls among preclinically disabled older people. A randomized controlled trial. *Am Med Dir Assoc*, 420-426. doi:https://doi.org/10.1016/j.jamda.2015.01.089.
- D'Silva, e. (2015). Impact Complications on Balance and Falls Contribution of the Vestibular System Physical . *Health and Sport*.
- Elipoulus. (2010). *Batasan Lanjut Usia* . Medan: Medan Merdeka.
- Fauzan, & Aldy. (2016). Patofisiologi Penuaan Pada lansia . *geriatric Journal*, 112-119.
- Fauzan; Aldhi. (2019). *Proses Degeneratif Pada Lansia*. Bogor: Sinar Edukasi.
- Huang, C., & Yang, T.-h. (2018, September 10). The benefits of Taichi Exercise on Balance Control in Elderly During Stair-to-Floor Transition. *International Society of Biomechanics in Sports*, 35-39.
- Huang, Y., & Liu, X. (2015). Improvement of balance control ability and flexibility in the elderly Tai Chi Chuan (TCC) practitioners: a systematic review and metaanalysis. *Arch Gerontol Geriatr*. doi:https://doi.org/10.1016/j.archger.2014.10.016 .
- Huang, Z., Feng, Y., Li, Y., Lv, C., Zheng , Y., & Cheng, Y. (2017). Systematic review and meta-analysis: tai Chi for preventing falls in older adults. *BMJ Open*. doi:https://doi.org/10.1136/bmjopen-2016-013661.
- Hwang. (2016). Balance and Risk Fall of Elderly. *Medicine and Science: Health Care Quality Unit* , 112.

- Jayanti. (2016). *Mengenal Usia Lanjut dan Perawatannya*. Jakarta: Salemba Medika.
- Kemenkes RI. (2016). *Penduduk Lansia*. Jakarta: Kemenkes RI.
- Kemenkes RI. (2019). *Kementerian Kesehatan Republik Indonesia: Analisis Lansia di Indonesia*. Pusat Data dan Informasi. Jakarta Selatan: KEMENKES RI.
- Kholifah, S. N. (2016). *Modul Bahan Ajar Keperawatan Gerontik Konsep Lanjut Usia dan Proses Penuaan*. Jakarta: kemenkes pusat pendidikan sumber daya manusia kesehatan.
- Kim, H. D. (2019). Effects of tai chi exercise on the center of pressure trace during obstacle crossing in older adults who are at a risk of falling. *Journal Physical Therapeutic Science*, 21(1), 49-54. doi:<https://doi.org/10.1589/jpts.21.49>
- Landi, F., Liperoti, R., Russo, A., Rosii, Q., Liper, & Forze. (2012). Sarcopenia as a risk factor for falls in elderly individuals: results from the iSIRENTE study. *Clinical Nutr*, 652-658. doi:<https://doi.org/10.1016/j.clnu.2012.02.007>.
- Maryam. (2010). *Pengaruh Resiko jatuh Tubuh Lansia*. Bandung: Medika Bandung.
- Miller, S., & Taylor-Piliae, R. (2014). Effects of Tai Chi on cognitive function in community-dwelling older adults: a review. *Geriatric Nurse*, 9-19. doi:<https://doi.org/10.1016/j.gerinurse.2013.10.013>.
- Nagai, K., Yamada, M., Uemura, Y., Yamada, Y., Ichihashi, N., & Tsuboyama, T. (2011). Differences in muscle coactivation during postural control between healthy older and young adults. *Archives of Gerontology and Geriatrics*. doi:<https://doi.org/10.1016/j.archger.2011.01.003>
- Nugroho. (2008). *Batasan Usia Lansia*.
- Nursalam. (2020). *Metodelogi Penelitian Ilmu Keperawatan*. Jakarta : Salemba Medika.
- Okuyan, C. B., & Deveci, E. (2020, Oktober 31). The effectiveness of Tai Chi Chuan on fear of movement, prevention of falls, physical activity, and cognitive status in older adults with mild cognitive impairment: A randomized controlled trial. *Perspect Psychiatr Care*. doi:DOI: 10.1111/ppc.12684
- Peterson, d., & Martin, P. (2010). Effects of age and walking speed on coactivation and cost of walking in healthy adults. *Gait and Posture*, 31(355), 11-15. doi:<https://doi.org/10.1016/j.gaitpost.2009.12.005>

- Savira. (2016). *Buku Ajar Keperawatan Medikal Bedah*. Jakarta: EGC.
- Sugiono. (2018). *Sumber data pencarian literature*. Jakarta: Jakarta Salemba.
- Suhartono. (2006). *Konsep Resiko jatuh Tubuh*. Jakarta: Jakarta Medika.
- Sun, W., Ma, X., Wang, L., Zhang, C., Song, Q., Gu, H., & Mao, D. (2018). Effects of Tai Chi Chuan and Brisk Walking Exercise on Balance Ability in Elderly Women: A Randomized Controlled Trial. *Human Kinetics Article*, 1-15. doi:<https://doi.org/10.1123/mc.2017-0055>
- United Nations. (2017). *World Population Prospect*. Chicago: United Nations int .
- Wang, H., Wei, A., Lu, Y., Chen, W., Chang, M., & Huang, A. (2016). Simplified tai chi program training versus traditional tai chi on the functional movement screening in older adults. *Evid-Based Complementary Altern Med*. doi:<https://doi.org/10.1155/2016/5867810>.
- WHO. (2018). *Elderly People*. Jenewa, Swiss.
- Widarti. (2020, April 9). Pengaruh Latihan Taichi terhadap Resiko jatuh Lansia. *Jurnal Medis Research*, 121.
- Wong, T. W. (2019, Mei 5). Feasibility and preliminary efficacy of Ai Chi aquatic exercise training in Hong Kong's older adults with risk of falling: Design and methodology of a randomized controlled trial. *Contemporary Clinical Trials Communications*, 100. doi:<https://doi.org/10.1016/j.conctc.2019.100376>
- World Health Organization. (2018, Agustus 23). *WHO International* . Retrieved from World Health Organization: <https://www.who.int/>

LAMPIRAN

Lampiran 1 Artikel

Journal of Exercise Science & Fitness 18 (2020) 142–147

Contents lists available at ScienceDirect

Journal of Exercise Science & Fitness

journal homepage: www.elsevier.com/locate/jest

Augmented reality-assisted training with selected Tai-Chi movements improves balance control and increases lower limb muscle strength in older adults: A prospective randomized trial

Po-Jung Chen^{a,1}, I-Wen Penn^{b,c,1}, Shun-Hwa Wei^a, Long-Ren Chuang^d, Wen-Hsu Sung^{a,*}

^a Department of Physical Therapy and Assistive Technology, National Yang-Ming University, Taipei, 11221, Taiwan

^b School of Medicine, Fu Jen Catholic University, New Taipei City, 24203, Taiwan

^c Department of Physical Medicine and Rehabilitation, Fu Jen Catholic University Hospital, Fu Jen Catholic University, New Taipei City, 24103, Taiwan

^d Department of Chinese Martial Arts, Chinese Culture University, Taipei, 11114, Taiwan

ARTICLE INFO

Article history:
Received 20 March 2020
Received in revised form 20 May 2020
Accepted 20 May 2020
Available online 21 May 2020

Keywords:
Tai-Chi
Exercise
Muscle strength
Functional balance
Augmented reality-assisted training

ABSTRACT

Background: Tai-Chi benefits older adults by enhancing balance control and increasing the muscle strength of the lower limbs. However, a complete set of traditional Tai-Chi exercises is sometimes too difficult for beginners. We investigated whether practicing augmented reality-assisted training with selected Tai-Chi movements tailored to the practitioner's ability (selected Tai-Chi, or sTC) is as effective as performing a complete set of Tai-Chi sequences (complete traditional Tai-Chi, or tTC).

Methods: In this prospective randomized trial carried out in the Beitou District of Taipei City, Taiwan, community-dwelling adults aged ≥ 65 and without any debilitating diseases ($n = 28$) were included. Participants were randomly assigned to the sTC group ($n = 14$) or the tTC group ($n = 14$). Participants in the sTC group practiced selected Tai-Chi movements using the augmented reality Tai-Chi training system. Participants of the tTC group were asked to complete the 24-form Yang-style Tai-Chi following the instructions of Tai-Chi masters. Each training session lasted 30 min, with 3 sessions per week for 8 weeks. Pre- and post-intervention evaluations included functional balance tests, comprising the Berg Balance Scale (BBS), Timed Up and Go test (TUG), and Functional Reach Test (FRT), as well as muscle strength measurements of the lower extremities.

Results: Pre-intervention evaluations showed significant differences in FRT ($p = 0.034$) and left hip abductor muscle strength ($p = 0.016$) between the sTC and tTC groups. After 8 weeks of training, the BBS, TUG, and FRT scores in the sTC group showed significant improvement overall. Although all three functional balance test scores improved in the tTC group, only the improvement in BBS was statistically significant ($p = 0.001$). After 8 weeks, all muscle strength measurements increased by an average of 3.1 ± 1.0 kgw in the sTC group and 1.5 ± 0.8 kgw in the tTC group.

Conclusions: The augmented reality-assisted training with selected Tai-Chi movements, designed based on objective measurements of the practitioner's capability, improved balance control and muscle strength of lower limbs at least as effectively as the complete sequence of traditional Tai-Chi exercises. Trial registration: This study was approved by the Institutional Review Board of National Yang-Ming University (IRB number: 1090087). Written informed consent was obtained from all participants.

© 2020 The Society of Chinese Scholars on Exercise Physiology and Fitness. Published by Elsevier (Singapore) Pte Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Abbreviations: augmented reality, AR; base of support, BOS; complementary metal-oxide-semiconductor, CMOS; Berg Balance Scale, BBS; center of pressure, COP; electrogoniometer, EHG; Functional Reach Test, FRT; selected Tai-Chi, sTC; Timed Up and Go test, TUG; traditional Tai-Chi, tTC; virtual reality, VR.

* Corresponding author.
E-mail addresses: whsung@ym.edu.tw, wenhsu.sung@gmail.com (W.-H. Sung).

Po-Jung Chen and I-Wen Penn have equally contributed to this article.

Introduction

Falls are a significant threat to the quality of life in older adults, often causing a decline in self-care and in the participation in physical and social activities. Falls have multifactorial causes. Prominent intrinsic fall risk factors are age-related decreases in

<https://doi.org/10.1016/j.jest.2020.05.005>

1728-8626/© 2020 The Society of Chinese Scholars on Exercise Physiology and Fitness. Published by Elsevier (Singapore) Pte Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

balance and strength/power performance, as well as cognitive decline.¹ In the elderly, loss of balance often results from sarcopenia and frailty,² and from their inability to shift their feet positions to allow adequate base of support (BOS) function.^{3,4} To prevent falls, various training programs have been developed to enhance functional balance control and increase the muscle strength of the lower extremities, with some tailored specifically for the elderly.^{5,6}

Tai Chi, characterized by a sequence of gentle, low-impact, and coordinated movements, is an appropriate form of exercise for older adults because it involves minimal strain on joints and the cardiovascular system.⁷ Previous studies have shown that Tai Chi increased muscle strength in the lower extremities, and improved balance control, proprioception, and postural adaptation, thereby reducing fall risk in older adults.^{8–11} However, since aging is an individualized process, prescription of exercises, such as Tai Chi, should be customized based on the health status and physical functioning of each individual. We have previously demonstrated that a set of simplified Tai Chi exercises, consisting of only three to five movements, selected based on the subjective measurements of the practitioner's capability, could be beneficial to people with no prior experience with Tai Chi.^{12,13}

Augmented Reality (AR) is a technology that projects virtual information on the real world through image recognition, which can increase users' awareness and understanding of their surroundings.¹⁴ In addition to games and entertainment, AR technology is currently used in medical-related fields, such as medical teaching and training, surgical simulation, and neurological rehabilitation.^{15,16} In 2017, Yoo et al. used AR with electromyography (EMG) signals for neuromuscular coordination training in children with cerebral palsy. Compared to using EMG signal feedback alone, EMG combined with AR resulted in better neuromuscular effects on elbow control.¹⁷ In the same year, Villiger et al. applied AR technology to assist in training lower limbs in patients with incomplete spinal cord injury and found significant improvements in lower limb muscle strength, balance, and functional activity.¹⁸ Using AR with the Origa Exercise has significantly improved the strength of muscles associated with knee flexion and ankle dorsiflexion in older women, and the scores of the Morse Fall Scale among the same participants were significantly improved. There was also an improvement in the degree of asymmetry relative to the body center of gravity that provides a measure of balance.¹⁹

In the present study, we enhanced the participants' performance of the simplified Tai Chi exercises by using an AR training system. We compared the beneficial effects of this AR-assisted selected Tai Chi (sTC) intervention with the complete sequence of traditional Tai Chi (TC) exercises.

Methods

This study was a prospective randomized trial. Participants received baseline assessments, with follow-up after 8 weeks. The study was approved by the Institutional Review Board of the National Yang-Ming University (#1000087). Written informed consent was obtained from all participants.

Participants

As shown in Fig. 1, 28 residents of the Beitou District of Taipei City, Taiwan participated in the study. The inclusion criteria required that participants were ≥ 65 years of age and achieved a score of ≥ 24 on the Mini-Mental Status Examination (MMSE), indicating normal cognitive function. A certified physical therapist performed the evaluation. Exclusion criteria were disorders of the central nervous system or musculoskeletal system, including strokes, head injuries, Parkinson's Disease, fractures, or the




Fig. 1. Flowchart explaining the assignment of the participants to the selected Tai Chi (sTC) group and the complete traditional Tai Chi (TC) group.

presence of orthopedic implants. Individuals with previous experience in Tai Chi exercise were also excluded. After assignment to the exercise groups, all participants completed the 8 weeks of sessions as advised.

The AR-assisted training system

The AR-assisted Tai Chi training system was developed to facilitate the practice of Tai Chi. Microsoft Kinect system (Microsoft Corporation) was used as a body motion detector. Kinect has a three-dimensional depth sensor composed of an infrared emitter and an infrared complementary metal-oxide semiconductor (CMOS) camera. This system can capture the movements of the entire body, as well as gestures and sounds. The Kinect software development kit was utilized to construct the digital skeleton from the spatial position of all four limbs of the performer and calculate skeletal information.²⁰ We used the skeletal tracking system to detect critical changes in movement of the lower limbs and collect movement data when a participant stepped forward or moved to the side. As shown in Fig. 2, a computer vision-based interaction method was employed where the skeletal data and the movements of the Tai Chi coach were displayed to create an AR scenario for the practitioner to follow. The movements and the skeletal information of participants were also displayed. Thus, the practitioners could compare their performance of the exercises with that of the




Fig. 2. Setup for the augmented reality-assisted Tai-Chi training system.

masters. The accuracy of the practitioner's movement using the skeletal information of the coach as a reference was calculated and the practitioner received real-time feedback.

Selected Tai-Chi (sTC) exercise

We monitored the base of support (BOS) of each participant by measuring the displacement of the Center of Pressure (COP) using a computerized evaluation system.^{12,13} We extracted 8 simple steps from the original 24-form Yang-style Tai-Chi sequences, based on the COP analyses, for development of computer-assisted Tai-Chi exercises. Participants in the sTC group were asked to perform COP tests and were assigned one individual Tai-Chi movement at an appropriate difficulty level based on their maximum COP displacement and endurance.¹³ Participants in the sTC group repeated the individually selected movement for 30 min in each training session, with three sessions per week for 8 weeks. The exercise program commenced with movements of low difficulty, then progressed to higher difficulty as participants became adept at each movement.^{13,20}

Traditional Tai-Chi (tTC) exercise

Participants of the complete traditional Tai-Chi (tTC) program learned and practiced the classical 24-form Yang-style Tai-Chi exercise under the instruction of a certified Tai-Chi master. The exercise program consisted of three sessions per week for 8 weeks and the duration of each session was 30 min. During each session, participants usually practiced a complete Yang-style Tai-Chi exercise two or three times.

Outcome measurements

Functional balance tests, including the Berg Balance Scale (BBS), Timed Up and Go test (TUG) and Functional Reach Test (FRT), as well as measurements of lower-extremity muscle strength were conducted before and after the Tai-Chi intervention. The tests and measurements were conducted by a certified physical therapist who was blinded to the sTC or tTC exercise allocation.

The BBS consisted of 14 items that evaluated the balance of each individual in sitting, standing, and transitional positions. The scores were determined using an ordinal 5-point scale with a total score ranging from 0 to 56, where higher scores indicated better balance control.²²

The TUG is a simple test used to assess a person's mobility that requires both static and dynamic balance.²³ The test measures the time taken by a participant to stand up from a chair, walk 3 m at a comfortable speed, turn around, return to the chair, and then sit down. Three timed trials are needed to ensure performance stability in the TUG test. Mean values for three trials, with a 1-min rest between each trial, were used for analysis.

The FRT was used to assess dynamic balance. This test measures the maximal forward distance a participant could reach beyond the length of their arms. Participants were asked to stand next to a wall and reach forward as far as possible without moving their feet. The extra reaching distance was then recorded in centimeters.²⁴

Lower-extremity muscle strength was measured using a handheld isometric dynamometer, MicroFET®3 (Hoggan Health Industries). This portable digital dynamometer performs accurate, objective muscle testing and enables assessors to stabilize and assist individuals while keeping one hand free. In this study, the following 8 lower-limb muscles were bilaterally measured: hip flexor, hip extensor, hip abductor, hip adductor, knee extensor, knee flexor, ankle dorsi flexor, and ankle plantar flexor. The participants were asked to perform maximal isometric contractions for 3 s. The

average of three measurements, at a 60-sec resting interval, were calculated and recorded.

Statistical analysis

Most of the parameters showed normal distribution and were evaluated with Shapiro-Wilk tests; only the TUG and left hip extensor muscle strength measurements of the sTC group and right ankle dorsi-flexor muscle strength measurements of tTC group showed non-normal distribution. If the parameters demonstrated normal distribution, we use a paired t-test for the within-group test and an independent t-test for the between-group test. If the parameters demonstrated non-normal distribution, the Wilcoxon test was used for the within-group test and the Mann-Whitney test was used for the between-group test. If the parameter used nonparametric statistics, it was indicated via "*" in the table. All statistical analyses were performed using PASW Statistics version 18.0 (SPSS Inc.). Statistical significance was defined as a p-value < 0.05.

Results

Demographic characteristics

A flowchart depicting the assignment of the participants to the sTC group and tTC group is shown in Fig. 1. The demographic characteristics of the participants are listed in Table 1. No significant difference was observed in the distribution of age, sex, height, and weight between the sTC and tTC groups.

Significant improvements in balance control and muscle strength were observed following the Tai-Chi interventions

The baseline and follow-up evaluation results in the sTC and tTC groups are presented in Table 2. The evaluations consisted of 3 functional balance tests, BBS, TUG, and FRT, and lower-limb muscle strength assessments. In the sTC group, BBS, TUG, and FRT scores showed significant improvement overall. The strength of each lower limb muscle also increased by an average of 3.1 ± 1.0 kgw. In the tTC group, although all three functional balance test scores improved, only the BBS improvement was statistically significant ($p = 0.001$). The muscle-strength assessments showed increases in all lower limb muscles, with an average of 1.6 ± 0.8 kgw. Improvements of the right hip flexor (RH F, $p = 0.032$), left hip flexor (LH F, $p = 0.033$), left hip abductor (LH Ab, $p = 0.001$), right ankle dorsi-flexor (RA Df, $p = 0.001$), and left ankle dorsi-flexor (LA Df, $p = 0.002$) reached statistical significance. Comparing the baseline and follow-up assessments in the sTC and tTC groups, participants in the sTC group exhibited improvements in all 19 items after the sTC intervention, while only 6 out of 19 items showed improvement after the tTC intervention.

Table 1
Demographic characteristics of participants.

	sTC Group	tTC Group	p value
Number	14	14	
Male	2	1	
Female	12	13	
Age, y/o, mean $\pm$ SD	72.2 $\pm$ 2.8	73.1 $\pm$ 5.5	0.501
Height, cm, mean $\pm$ SD	156.1 $\pm$ 6.1	156.2 $\pm$ 3.8	0.449
Weight, kg, mean $\pm$ SD	59.1 $\pm$ 8.6	58.1 $\pm$ 6.6	0.775

sTC = selected Tai-Chi; tTC = traditional Tai-Chi; SD = standard deviation.

Table 2
Baseline and follow-up evaluation results in sTC and tTC groups.

	sTC Group (n = 14)			tTC Group (n = 14)		
	Baseline (mean ± SD)	Follow up (mean ± SD)	p value	Baseline (mean ± SD)	Follow up (mean ± SD)	p value
Functional Balance						
BBS	50.3 ± 2.1	54.0 ± 1.1	<0.001*	49.2 ± 4.5	51.1 ± 4.7	0.001*
TUG	8.7 ± 0.7	6.9 ± 0.9	<0.001*	9.0 ± 1.8	8.4 ± 1.6	0.115
FRT	25.7 ± 4.2	32.5 ± 4.1	<0.001*	24.3 ± 5.6	24.8 ± 3.8	0.788
Muscle Strength						
RH F	17.7 ± 6.1	23.7 ± 7.5	0.001*	16.6 ± 7.9	18.5 ± 5.1	0.002*
LH F	17.1 ± 6.2	23.7 ± 7.7	0.001*	15.4 ± 4.3	17.7 ± 5.4	0.003*
RH E	17.1 ± 5.1	20.8 ± 7.4	0.007*	16.4 ± 3.9	18.2 ± 3.8	0.001*
LH E	17.8 ± 5.8	21.5 ± 8.6	0.001*	16.2 ± 2.9	17.1 ± 5.4	0.310
RH Ab	12.9 ± 3.4	15.8 ± 5.9	0.003*	15.7 ± 3.4	16.8 ± 2.6	0.276
LH Ab	12.5 ± 3.2	15.7 ± 5.5	0.002*	17.1 ± 7.0	15.8 ± 2.9	0.002*
RH Ad	11.6 ± 3.2	14.3 ± 4.8	0.005*	11.1 ± 2.1	12.9 ± 2.9	0.090
LH Ad	10.7 ± 3.2	13.9 ± 4.7	0.007*	12.1 ± 2.8	13.4 ± 2.6	0.121
RH K	11.7 ± 3.7	17.7 ± 4.0	0.010*	17.1 ± 2.6	14.1 ± 2.1	0.573
LH K	11.0 ± 3.7	13.1 ± 4.9	0.009*	12.8 ± 2.6	13.8 ± 2.9	0.226
RH F	10.1 ± 4.2	21.4 ± 5.8	0.010*	18.2 ± 3.5	18.8 ± 4.0	0.478
LH F	18.1 ± 4.0	21.2 ± 5.6	<0.001*	16.9 ± 2.4	17.8 ± 4.1	0.484
RA DF	13.3 ± 3.9	15.4 ± 5.0	0.003*	11.1 ± 2.3	13.9 ± 2.7	0.001*
LA DF	12.5 ± 3.7	15.1 ± 4.7	<0.001*	11.4 ± 2.5	14.0 ± 2.9	0.002*
RA PF	18.3 ± 4.1	22.8 ± 8.3	0.015*	17.8 ± 2.5	19.3 ± 3.6	0.128
LA PF	18.6 ± 3.9	21.7 ± 6.7	0.010*	16.7 ± 2.8	17.3 ± 4.0	0.446

* nonparametric statistics; *p < 0.05; sTC = selected Tai-Chi; tTC = traditional Tai-Chi; SD = standard deviation; BBS = Berg Balance Scale; TUG = Timed Up and Go test; FRT = Functional Reach Test; RH = right hip; F = flexor; LH = left hip; E = extensor; Ab = abductor; Ad = adductor; RK = right knee; LK = left knee; RA = right ankle; DF = dorsiflexor; LA = left ankle; PF = plantar flexor.

Beneficial effects of the AR-assisted selected Tai-Chi program were comparable to the complete Tai-Chi sequence.

Comparisons of the functional balance tests and muscle strength measurements between the sTC and tTC groups are shown in Table 3. Pre-intervention evaluations showed significant differences in FRT ($p = 0.034$) and RH Ab ($p = 0.046$) between the sTC and tTC groups and post-intervention evaluations showed significant differences in BBS ($p = 0.044$), TUG ($p = 0.015$), and FRT ($p < 0.001$) between groups.

Discussion

The Kinect depth sensor and skeleton tracking system senses the body movement of the participant without using other input devices and limits the disturbance of movement, which is especially important for elderly people.

The measurements used in the current study were not tests specific to Tai-Chi, however, assessments such as BBS^{25,26}, TUG²⁷, and lower limb muscle strength^{28–30} have been widely used in the other Tai-Chi studies. In this study, we found that both sTC and tTC

Table 3
Baseline and follow-up evaluation results between sTC and tTC groups.

	Baseline			Follow up		
	sTC group (mean ± SD)	tTC group (mean ± SD)	p value	sTC group (mean ± SD)	tTC group (mean ± SD)	p value
Functional Balance						
BBS	50.3 ± 2.1	49.2 ± 4.5	0.444	54.0 ± 1.1	51.1 ± 4.7	0.044*
TUG	8.7 ± 0.7	9.0 ± 1.8	0.390*	6.9 ± 0.9	8.4 ± 1.6	0.015*
FRT	25.7 ± 4.2	24.3 ± 5.6	0.034*	32.5 ± 4.1	24.8 ± 3.8	<0.001*
Muscle Strength						
RH F	17.7 ± 6.1	16.6 ± 7.9	0.394	23.7 ± 7.5	18.5 ± 5.1	0.115
LH F	17.1 ± 6.2	15.4 ± 4.3	0.411	23.7 ± 7.7	17.7 ± 5.4	0.094
RH E	17.1 ± 5.1	16.4 ± 3.9	0.573	20.8 ± 7.4	18.2 ± 3.8	0.283
LH E	17.8 ± 5.8	16.2 ± 2.9	0.390*	21.5 ± 8.6	17.1 ± 5.4	0.301*
RH Ab	12.9 ± 3.4	15.7 ± 3.4	0.046*	15.8 ± 5.9	16.8 ± 2.6	0.580
LH Ab	12.5 ± 3.2	12.5 ± 2.9	0.947	15.7 ± 5.5	15.8 ± 2.9	0.926
RH Ad	11.6 ± 3.2	11.1 ± 2.1	0.543	14.3 ± 4.8	12.9 ± 2.9	0.372
LH Ad	10.7 ± 3.2	12.1 ± 2.8	0.178	13.9 ± 4.7	13.4 ± 2.6	0.790
RH K	11.7 ± 3.7	13.3 ± 2.6	0.207	17.7 ± 4.0	14.1 ± 2.5	0.320
LH K	11.0 ± 3.7	12.8 ± 2.6	0.151	13.1 ± 4.9	13.8 ± 2.9	0.573
RH F	10.1 ± 4.2	18.2 ± 3.5	0.351	21.4 ± 5.8	18.8 ± 4.0	0.234
LH F	18.1 ± 4.0	16.9 ± 2.4	0.382	21.2 ± 5.6	17.8 ± 4.1	0.287
RA DF	13.3 ± 3.9	11.1 ± 2.3	0.089*	15.4 ± 5.0	13.9 ± 2.7	0.476*
LA DF	12.5 ± 3.7	11.4 ± 2.5	0.416	15.1 ± 4.7	14.0 ± 2.9	0.468
RA PF	18.3 ± 4.1	17.8 ± 2.5	0.582	22.8 ± 8.3	19.3 ± 3.6	0.171
LA PF	18.6 ± 3.9	16.7 ± 2.8	0.380	21.7 ± 6.7	17.3 ± 4.0	0.064

* nonparametric statistics; *p < 0.05; sTC = selected Tai-Chi; tTC = traditional Tai-Chi; SD = standard deviation; BBS = Berg Balance Scale; TUG = Timed Up and Go test; FRT = Functional Reach Test; RH = right hip; F = flexor; LH = left hip; E = extensor; Ab = abductor; Ad = adductor; RK = right knee; LK = left knee; RA = right ankle; DF = dorsiflexor; LA = left ankle; PF = plantar flexor.

training could improve a practitioner's balance control and increase lower limb muscle strength after an 8-week intervention. Furthermore, the added benefits of performing simplified and personalized Tai-Chi exercises were substantial compared to the complete set of traditional Tai-Chi sequences, even if not to the level of statistical significance. An explanation of these results might be that the traditional Tai-Chi sequences contained movements that were too difficult for elderly practitioners. It has been shown that the amplitude of weight-shifting estimated by the degree of COP displacements differed among Tai-Chi movements.³¹ Practitioners with limited ability to voluntarily shift their weight in different spatial directions and to briefly maintain stability in different positions may encounter difficulty in conducting the movements. Thus, not all older individuals can complete the full set of exercises included in a traditional Tai-Chi program. We noted that the participants in the RTC group often omitted or skipped complicated movements or performed the steps at their own will. Therefore, older participants may require special considerations when a Tai-Chi exercise program is developed for them.³²

Many researchers have tried to simplify Tai-Chi programs.^{33–36} However, most of the simplified exercises were designed primarily based on the advice of experts rather than on the practitioner's abilities, and the training course was typically conducted at a pre-determined level of difficulty. With such a design, the only option for many participants would be to give up when encountering movements that were beyond their capabilities. For example, the one-legged station step of the traditional Tai-Chi exercise was too difficult for most elderly practitioners, yet could be readily accomplished by some older individuals. Therefore, simplified Tai-Chi exercises need to be customized to achieve optimal outcomes. We previously reported the development of personalized Tai-Chi exercise programs for older adults based on the practitioners' COP displacement values.^{12,13} Therefore, the individual Tai-Chi movements assigned to the practitioner were not only feasible but were also at an appropriate level of difficulty. In the present study, we have developed this sTC program further by incorporating virtual reality (VR)-augmented or AR-assisted Tai-Chi training systems²⁷ in the exercise intervention protocols. The trainees could perform individual Tai-Chi movements at a high level of accuracy by comparing their own skeletal information with that of the Tai-Chi master, thereby achieving increased optimal training goals set by the trainer.

In the present study, participants in the sTC group were asked to practice only one movement repeatedly in each session. The movement was assigned according to the practitioner's balance control as assessed through COP measurements. The training was also designed to be progressive, with the practitioner continuously challenged in a gradual manner. When the original movement was learned and mastered, the trainer assigned another Tai-Chi movement that required an increased COP displacement or was at a higher difficulty level. The superior training results observed in the sTC group might have been attributable to the individually tailored Tai-Chi movements and the progressive intensity and complexity of the practice.

Tai-Chi exercises require the coordination of breathing and motor functions. It has been reported that Tai-Chi movements stimulated the part of the brain that governs balance, thereby increasing overall steadiness.^{35–41} Previous studies have also shown that a relatively long period of practice, typically from 12 to 48 weeks, was required for older practitioners to benefit from Tai-Chi exercises.^{3,35,42,43} In our study, we demonstrated that merely 8 weeks of intervention with simplified and personalized Tai-Chi training could provide significant benefit, comparable to that achieved with extended periods of Tai-Chi practice. Thus, simplified and personalized Tai-Chi exercises are particularly useful for Tai-Chi beginners and provide substantial benefit for those who

cannot learn and practice the complete Tai-Chi sequence, such as frail or disabled elderly individuals and those who cannot comprehend or memorize the full Tai-Chi sequence. Moreover, the AR-assisted Tai-Chi training program developed in the present study can serve as an operative assistive methodology to help the practitioner exercise without the physical presence of a coach, improve exercise adherence, achieve the trainer's goals, and maximize the training outcome. The results of our study suggest that performing selected Tai-Chi exercises was at least as useful and, for certain practitioners, even more beneficial than conducting the full sets of traditional Tai-Chi exercises.

The primary limitations of this study are the relatively small number of participants and imbalance among the sexes. There were more female individuals recruited than males. It is not possible to separate the statistics based on sex. Thus, the conclusions from this study should be made with caution, with the caveat that the evidence supports the benefit of sTC only for women. The included participants had prior exercise habits, but none had previous experience with Tai-Chi. The frequency of exercise of the participants was generally 2–3 times per week and types of exercise included walking, swimming, and social dance. The study did not ask participants to stop their current exercise during the intervention. In addition, the functional assessments of this study focused mainly on balance control and muscle strength of the lower limbs, without considering other beneficial effects typically associated with the complete set of traditional Tai-Chi sequences. It would be informative to conduct a follow-up survey to compare differences in cognition, adherence rate, or personal mental health status between the RTC and sTC group participants in the future.

Conclusions

We found that simplified and personalized Tai-Chi training, designed based on objective measurements of the practitioner's capability and conducted under the guidance of a VR-augmented training system with progressive intensity and complexity, could achieve training goals more readily than use of traditional full-course Tai-Chi, especially for beginners or laypersons not familiar with this type of exercise.

Funding

This study was supported by the Ministry of Science and Technology of Taiwan (NSC 100-2410-H-010-010-MY3). The funder did not have a role in conducting this study nor in writing the manuscript.

Declaration of competing interest

The authors declare that there is no conflict of interest.

CRediT authorship contribution statement

Po-Jung Chen: Conceptualization, Methodology, Software, Investigation, Data curation, Writing – original draft. **I-Wen Penn:** Conceptualization, Formal analysis, Resources, Writing – review & editing. **Shun-Hwa Wei:** Methodology, Supervision. **Long-Ren Chuang:** Methodology, Resources, Supervision. **Wen-Hsu Sung:** Conceptualization, Methodology, Software, Validation, Visualization, Supervision, Project administration.

Acknowledgements

The authors would like to express their gratitude to the Department of Physical Therapy and Assistive Technology, National

Yang-Ming University, for the wonderful support during the research.

References

1. Kim H.D. Effects of tai chi exercise on the center of pressure trace during obstacle crossing in older adults who are at a risk of falling. *J Phys Ther Sci*. 2019;21(11):40–44. <https://doi.org/10.1589/jpts.41.40>.
2. Landi F, Liperoti R, Russo A, et al. Sarcopenia as a risk factor for falls in elderly individuals: results from the DISABEST study. *Age Ageing*. 2012;41(5):652–658. <https://doi.org/10.1093/agea/afz202>.
3. Hillard MJ, Martinez KM, Janssen I, et al. Lateral balance factors predict future falls in community-living older adults. *Arch Phys Med Rehabil*. 2008;89(9):1704–1711. <https://doi.org/10.1016/j.apmr.2008.06.015>.
4. Nakamura H, Tsuchida T, Mizuo Y. The assessment of posture control in the elderly using the displacement of the center of pressure after forward platform translation. *J Electromyogr Kinesiol*. 2001;11(6):395–403. [https://doi.org/10.1016/S1550-5687\(01\)00018-5](https://doi.org/10.1016/S1550-5687(01)00018-5).
5. Gichwindi Y, Kessig BW, Lacroix A, Muehlbauer T, Heninger B, Granacher U. A best practice fall prevention exercise program to improve balance, strength, power, and psychosocial health in older adults: study protocol for a randomized controlled trial. *BMC Geriatr*. 2018;18:105. <https://doi.org/10.1186/s14714-018-0115-1>.
6. M. R. Exercise prescription for healthy populations with special considerations. In: *Medicine & Sport: An ACSM's Guidelines for Exercise Testing and Prescription* (edn edn). Philadelphia: Wolters Kluwer/Lippincott Williams & Wilkins Health; 2018:180–202.
7. Huang Y, Liu X. Improvement of balance control ability and flexibility in the elderly Tai Chi Chuan (TCC) practitioners: a systematic review and meta-analysis. *Arch Geriatr Geriatr*. 2015;50(2):231–238. <https://doi.org/10.1016/j.archger.2014.10.018>.
8. Huang ZG, Peng YH, Li YH, et al. Systematic review and meta-analysis: tai chi for preventing falls in older adults. *BMJ Open*. 2017;7(1):e013667. <https://doi.org/10.1136/bmjopen-2016-013667>.
9. Jin A, Taylor J, Zeng C, Song P. The effect of tai chi on functional lower extremity mobility and strength, ankle proprioception, and postural adaptation in older adults. *Am J Med Sci*. 2017;7:238–270. <https://doi.org/10.1001/jam.2017.070601>.
10. Song QH, Zhang QH, Xu RM, et al. Effect of Tai-chi exercises on lower limb muscle strength, bone mineral density and balance function of elderly women. *Int J Clin Exp Med*. 2014;7(9):1569–1578. <https://doi.org/10.1016/j.ijcem.2014.05.025>.
11. Tsang WW, Hui Chan CW. Effect of 4- and 8-week intensive Tai Chi Training on balance control in the elderly. *Med Sci Sports Exerc*. 2004;36(4):648–657. <https://doi.org/10.1249/01.mss.0000121541.57860.0f>.
12. Penn BW, Sung WH, Lin LH, Huang K, Huang Y, Lin PH. Effects of individualized Tai-Chi on balance and lower-limb strength in older adults. *BMC Geriatr*. 2019;19(1):215. <https://doi.org/10.1186/s12877-019-1230-5>.
13. Song WH, Lin CC, Wei SH, et al. Feasibility and outcome of an individualized Tai Chi program for improving balance and strength in the elderly: a pilot study. *NeuroRehabil*. 2018;33(4):500–515. <https://doi.org/10.1080/10791822.2018.1502561>.
14. Kamachi A, Kato S, Cho ST, et al. Application of virtual, augmented, and mixed reality to neurology. *Int J Neurosci*. 2016;20(3):172–181. <https://doi.org/10.1080/10791822.2016.1163577>.
15. Varea P, Roman J, Zanca P, et al. Recent development of augmented reality in surgery: a review. *J Health Eng*. 2017;2017:4534172. <https://doi.org/10.1155/2017/4534172>.
16. Valero-Cabré C, Borra L, Vignaroli-García F, Merello M. Development of a haptic interface for motor rehabilitation therapy using augmented reality. *Int Conf Proc IEEE Eng Med Biol Soc*. 2015. 2015:1156–1159. <https://doi.org/10.1109/EMBC.2015.7318171>.
17. Yoo JW, Lee DR, Cha YJ, You SH. Augmented effects of EMG biofeedback method with virtual reality on neuromuscular control and movement coordination during reaching in children with cerebral palsy. *NeuroRehabil*. 2017;42(2):175–185. <https://doi.org/10.1080/10791822.2017.1301402>.
18. Villiger M, Linero J, Awa L, et al. Home-based virtual reality-augmented training improves lower limb muscle strength, balance, and functional mobility following chronic incomplete spinal cord injury. *Front Neurosci*. 2017;11:555. <https://doi.org/10.3389/fn.2017.00555>.
19. Lee J, Yoo HK, Lee HJ. Effects of augmented reality-based Taiji exercise on balance, gait, and physical fitness in elderly women to prevent falls: a randomized controlled trial. *J Phys Ther Sci*. 2017;29(9):1585–1589. <https://doi.org/10.1589/jpts.2017.1585>.
20. Kinect for windows. Microsoft. Accessed <https://developer.microsoft.com/en-us/windows/kinect>. Accessed March 18, 2019.
21. Juras C, Rajan S, Freddi A, Sobott G, Rodi B. Evaluation of the limits of stability (LOS) balance test. *J Hum Kinet*. 2008;19:51–62. <https://doi.org/10.2478/s11832-008-9011-0>.
22. Berg K, Wood-Dauphinee S, Williams JL. The Balance Scale: reliability assessment with elderly residents and patients with an acute stroke. *Scand J Rehabil Med*. 1995;27(1):23–36. <https://doi.org/10.1080/165019795433010>.
23. Podsiadlo D, Richardson S. The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc*. 1991;39(2):142–148. <https://doi.org/10.1111/j.1532-5415.1991.tb01616.x>.
24. Duncan PW, Weiner DK, Chandler J, Studenski S. Functional reach: a new clinical measure of balance. *J Geriatr*. 1990;35(5):M100–M102. <https://doi.org/10.1001/jgeriatr.1990.04500100>.
25. Dytter S, Karagözlü H, Durmaz B, Bilingir U, Çakır S. A randomized controlled trial of Turkish folk dance on the physical performance, balance, depression and quality of life in older women. *Arch Geriatr Geriatr*. 2009;48(1):84–88. <https://doi.org/10.1016/j.archger.2007.10.008>.
26. Logghe BA, Zeeuw FE, Verhagen AP, et al. Lack of effect of Tai Chi Chuan in preventing falls in elderly people living at home: a randomized clinical trial. *J Am Geriatr Soc*. 2009;57(1):70–75. <https://doi.org/10.1111/j.1532-5415.2008.02064.x>.
27. Dyer R, Schmittal S, Komarskaya I, Pechen R. Tai chi and low impact exercise effects on the physical functioning and psychological well-being of older people. *J Appl Gerontol*. 2007;36(4):431–451. <https://doi.org/10.1177/0275460707318015>.
28. Chen S, Lu T, Hong Y. Kinematic and electromyographic analysis of the push movement in tai chi. *Br J Sports Med*. 2003;37:339–344. <https://doi.org/10.1136/bjsm.37.3.339>.
29. Hsu CJ, Gregor RJ, Waddell DG, et al. The influence of Tai Chi training on the center of pressure trajectory during gait initiation in older adults. *Arch Phys Med Rehabil*. 2004;85(10):1593–1598. <https://doi.org/10.1016/j.apmr.2004.05.001>.
30. Wolfson L, Whipple K, Derby C, et al. Balance and strength training in older adults: intervention goals and Tai Chi maintenance. *J Am Geriatr Soc*. 1990;38(5):580–590. <https://doi.org/10.1111/j.1532-5415.1990.tb01713.x>.
31. Liang GS, Tang WM. Comparison of the linear characteristics of standing and strong Tai Chi forms. *Obstet Reprod*. 2005;30(25):1891–1900. [https://doi.org/10.1002/1552-008X\(200512\)30:25%3C1891::AID-OBST1891%3E3.0.CO;2](https://doi.org/10.1002/1552-008X(200512)30:25%3C1891::AID-OBST1891%3E3.0.CO;2).
32. Lemay JF, Gagnon DH, Nadeau S, Grangeon M, Gauthier C, Dugas C. Center of pressure total trajectory length is a complementary measure to maximum excursion to better differentiate multidirectional standing limits of stability between individuals with incomplete spinal cord injury and able-bodied individuals. *J NeuroEng Rehabil*. 2014;11:8. <https://doi.org/10.1186/1743-0003-11-8>.
33. Chen HA, Chen WT, Huang MH. Development of the simplified Tai Chi exercise program (S2P) for frail older adults. *Comput Ther Med*. 2006;14(1):250–265. <https://doi.org/10.1016/j.cmt.2005.05.007>.
34. Li FZ, Fisher K, Hunter P, Shira M. A simpler eight-form easy tai chi for elderly adults. *J Aging Phys Act*. 2003;11:206–218. <https://doi.org/10.1177/1063426903011002>.
35. Toussaint M, Corbeil H, Roy PM, Desrosiers J, Dubuc N, Hebert R. Efficacy of supervised Tai Chi exercises versus conventional physical therapy exercises in fall prevention for frail older adults: a randomized controlled trial. *Obstet Reprod*. 2013;35(17):1429–1435. [https://doi.org/10.1002/1552-008X\(201307\)35:17%3C1429::AID-OBST1429%3E3.0.CO;2](https://doi.org/10.1002/1552-008X(201307)35:17%3C1429::AID-OBST1429%3E3.0.CO;2).
36. Wang R, Wei A, Lu Y, et al. Simplified tai chi program training versus traditional tai chi on the functional movement screening in older adults. *Dual-based Complementary Altern Med*. 2018;2018:5857810. <https://doi.org/10.1155/2018/5857810>.
37. Hsieh CC, Lin PS, Hsu WC, et al. The effectiveness of a virtual reality based tai chi exercise on cognitive and physical function in older adults with cognitive impairment. *Demer Geriatr Cognit Disord*. 2018;45(6):358–370. <https://doi.org/10.1016/j.dem.2018.04.009>.
38. Li F, Hunter P, Fitzgerald K, et al. Tai chi and postural stability in patients with Parkinson's disease. *Park J Med*. 2012;46(8):511–515. <https://doi.org/10.1056/NEJMed1079111>.
39. Miller MA, Taylor-Pleat R. Effects of Tai Chi on cognitive function in community-dwelling older adults: a review. *Geriatr Nurs*. 2014;35(1):16–19. <https://doi.org/10.1016/j.gernu.2013.10.015>.
40. Wayne PM, Walsh JM, Taylor-Pleat R, et al. Effect of tai chi on cognitive performance in older adults: systematic review and meta-analysis. *J Am Geriatr Soc*. 2014;62(1):25–39. <https://doi.org/10.1111/jags.12611>.
41. Zheng C, Liu F, Li S, Huang M, Tao J, Chen L. Tai chi and the protection of cognitive ability: a systematic review of prospective studies in healthy adults. *Am J Prev Med*. 2015;49(1):80–97. <https://doi.org/10.1016/j.amepre.2015.01.002>.
42. Day L, Hill KB, Sathakis VZ, et al. Impact of tai-chi on falls among predominantly disabled older people: A randomized controlled trial. *J Am Med Assoc*. 2019;321(14):1401–1410. <https://doi.org/10.1001/jama.2018.181188>.
43. Taylor-Pleat R, Hunter PM, Hepworth JT, Lee LM, Nayak U, Cook RM. Effect of Tai Chi on physical function, fall rates and quality of life among older stroke survivors. *Arch Phys Med Rehabil*. 2014;95(5):815–824. <https://doi.org/10.1016/j.apmr.2014.05.001>.



Feasibility and preliminary efficacy of Ai Chi aquatic exercise training in Hong Kong's older adults with risk of falling: Design and methodology of a randomized controlled trial

Thomson W.L. Wong^a

^a School of Public Health, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, China

ARTICLE INFO

Keywords:
Fall prevention
Older adults
Ai Chi aquatic exercise
Land-based exercise
Feasibility
Efficacy

ABSTRACT

Falls in older adults are a major global public health concern. Group exercise could mitigate fall risk but the traditional land-based group exercise may not be always suitable for older adults at risk of falling, especially for those with musculoskeletal problems. Ai Chi aquatic exercise program could provide a safe and low-impact exercise training for older adults. However, the feasibility and efficacy of the program has not been well-investigated. The objective of this study is to investigate the feasibility and efficacy of the Ai Chi aquatic exercise program, compared to the land-based exercise program for older adults with moderate to high risk of falling. Forty community-dwelling older adults aged 65 or above with moderate to high fall risk will be recruited. They will be randomly allocated to the Ai Chi Aquatic Exercise Group (intervention) or the Land-based Exercise Group (active control) receiving 16 sessions (8 weeks) of specific exercise training. Feasibility of both exercise groups will be examined by recruitment, adherence, retention, feedback, subjective exercise experiences and satisfaction. Preliminary efficacy will be determined by whether physical and psychological fall risk factors could be mitigated. Physical fall risk assessment will include tests for flexibility, muscle strength, gait and balance. Psychological fall risk will be assessed by preliminary cognitive function, anxiety level, level of depression and fear of falling. The results could establish a solid foundation for worldwide development of a feasible, safe and effective Ai Chi aquatic exercise program for prevention of falls in older adults with risk of falling.

1. Introduction

With global aging, fall-related injuries among older adults have been increased continuously, stressing our healthcare systems [1]. A previous study revealed that the prevalence of falls and recurrent falls among Hong Kong older adults was 19.3% and 4.75%, respectively [2]. It is not uncommon that falls lead to undesirable physical and psychological consequences in older adults, such as immobility and fear of falling [3]. In severe incidences, falls may even result in deaths. Therefore, falling has been a prominent health-related problem in the older population in Hong Kong. In 2017, for example, there were 277 registered deaths and 44150 in-patient discharges and deaths in Hong Kong due to falls [4]. The fallers inevitably require additional resources to cure their fall-related health problems. As a result, falling in older adults imposes an extremely heavy economic burden to not only older fallers themselves but also the worldwide health care system. With the frightening trend towards a progressively increasing aging population

[5], these problems are likely to affect the society more comprehensively.

It is widely agreed that "prevention is better than cure". Therefore, prevention of falls in older adults is of ultimate importance. According to a recent systematic review, one of the most effective interventions for fall prevention is group exercise [6]. Very strong evidence had illustrated that performing a well-structured group exercise program regularly by older adults could reduce the risk and rate of falls [6]. The favourable influences of group exercise program as a single fall prevention intervention had been demonstrated to be comparable to those of multi-faceted interventions [6]. It has also been suggested that exercise components essential in preventing falls of older adults include strength, balance, flexibility and reaction time [7]. Nevertheless, owing to aging-related physiological changes, such as loss of bone density and diminishing muscular mass [8], land-based exercises are not always be suitable for those older adults who have high risk of falling due to the fear issues [9]. In order to address such influential limitations of land-

^a School of Public Health, Li Ka Shing Faculty of Medicine, The University of Hong Kong, 3/F, The Hong Kong Jockey Club Building for Interdisciplinary Research, 5 Sassoon Road, Pokfulam, Hong Kong SAR, China.
E-mail address: wongthl@pku.hk.

<https://doi.org/10.1016/j.conctc.2019.100376>

Received 23 November 2018; Received in revised form 6 March 2019; Accepted 5 May 2019

Available online 05 May 2019

2451-8654/ © 2019 The Author. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

based fall prevention group exercise program, aquatic group exercise program could potentially provide the effective components (i.e., strength and balance training) of fall prevention [7], while minimizing potential problematic issues (e.g., increasing joint pain and fear of falling). Aquatic group exercise program may provide an alternative training to land-based exercise for those older adults who have degenerative joint problems and fear of falling. It may also result in better fall prevention and rehabilitation outcomes.

The characteristics of water (e.g., buoyancy & viscosity) could be utilized in aquatic exercise to provide a 'cushion' environment by reducing loads on weight-bearing joints but at the same time deliver graded levels of resistance for older adults during exercise [30]. Based on these characteristics, aquatic exercise could provide a potentially feasible, safe and effective therapeutic environment for older adults to perform exercise. Aquatic exercise is indeed well proven, with its benefits in promoting balance and functional mobility. For instance, Lord et al. [10] reported that older adults who completed an aquatic exercise program for nine sessions gained significant improvements in lower limb muscle strength, postural sway, flexibility and joint pain compared with the control group. Comparable results were also found by Tancoulon et al. [11] showing that a 24-week aquatic training program of older adults improved muscular strength. Participants were found to have improvements in muscular strength, flexibility, trunk flexion as well as functional mobility. Providing such benefits, different aquatic exercises, for example, water aerobic exercises, water walking and water dancing had been developed in current fitness program for health improvement and rehabilitation of both healthy older adults and those with disabilities.

One potentially feasible, safe and effective aquatic exercise program for fall prevention in older adults is called Ai Chi, which involves slow-and-based movements of the limbs and body (Tai Chi element) in a continuously flowing pattern and deep breathing (Qi Gong element) [12]. Ai Chi involves the 19-style aquatic exercise program of 1, Contemplating, 2, Floating, 3, Uplifting, 4, Enclosing, 5, Folding, 6, Soothing, 7, Gathering, 8, Freeing, 9, Shifting, 10, Accepting, 11, Accepting with grace, 12, Rounding, 13, Balancing, 14, Flowing, 15, Reflecting, 16, Suspending, 17, Encircling, 18, Surrounding, and 19, Nurturing [12]. Despite the relatively small number of previous studies, pilot research had demonstrated its potential effectiveness in improving postural balance control for older adults with high and medium risk of falling [13]. Being an aquatic exercise program, Ai Chi provides a safe and low-impact training environment that has been well-recognized as a desirable exercise training environment for older adults [9]. Furthermore, Ai Chi has potential benefits beyond those associated with traditional aquatic exercise. The rationale is that Ai Chi's oriental roots echo well with exercises such as Tai Chi and Qi Gong, which Hong Kong older adults are likely to engage in. Therefore, Ai Chi may be considered as a potentially feasible, safe and effective fall-prevention exercise program for older adults in Hong Kong.

According to a recent review of water-based Tai Chi, there was scanty number of studies investigating on the effect of Ai Chi in reduction of fall risk factors for older adults [14]. The Ai Chi program conducted in previous pilot research was varied and may have not been designed by a group of qualified Ai Chi and rehabilitation experts [13]. In order to enhance the beneficial effects of Ai Chi in fall prevention, an Ai Chi aquatic exercise program for fall prevention in Hong Kong has to be developed by a group of Ai Chi and rehabilitation experts in Hong Kong. To examine the feasibility and preliminary efficacy of this developed Ai Chi aquatic exercise program, the feasibility and preliminary efficacy of this developed program should be compared to the traditional land-based exercise program.

This study protocol aims to investigate the feasibility and preliminary efficacy of the Ai Chi aquatic exercise program that was developed by a group of Ai Chi and rehabilitation experts in Hong Kong through a randomized controlled trial. With reference to the past pilot research, the Ai Chi aquatic exercise program may be more feasible and

effective than the traditional land-based fall prevention exercise program [13]. It is hypothesized that the expertly-designed Ai Chi aquatic exercise program could be as feasible as the land-based exercise program in the context of recruitment, adherence, retention and participants feedback, subjective exercise experiences and satisfaction of participants. It is also hypothesized that the expertly-designed Ai Chi aquatic exercise program is more effective in mitigating physical and psychological fall risk factors of Hong Kong's older adults with medium to high risk of falling, compared to the land-based exercise program.

2. Method

2.1. Study design

This study will be a randomized controlled trial. Eligible participants will be randomly assigned by concealed block randomization by an independent person with a sealed and opaque envelope system to either the Ai Chi Aquatic Exercise Group (intervention) or the Land-based Exercise Group (active control group). All participants will undergo two assessment sessions before training at baseline (T0) and just after completion of all exercise sessions (T1). In the baseline assessment (T0), a structured questionnaire will be used to ask for demographics, detailed medical history, detailed history of fall incident, social history and socio-economic status of the participants. During the two assessment sessions (T0 and T1), physical and psychological fall risk factors will be assessed. Feasibility outcome variables will be measured during and at the end of the whole study.

2.2. Participants

Forty older adults with moderate to high risk of falling will be recruited from different elderly community centres in Hong Kong by convenience sampling. Participants will only be included if they are aged 65 or above, score a total score of equal or more than 24 in the Chinese version of the Mini-Mental State Examination (MMSE-C) [15], score less than 24/28 in the Tinetti Balance Assessment Tool [16], have no history of any neurological diseases, and have no contraindication for aquatic exercises. The sample size was estimated based on a pilot study conducted by Teixeira and colleagues to compare the effect of a 6-week Ai Chi exercise group with control group on balance ability of older adults, an effect size of ≈ 1.1 of the primary outcome (balance score of the Tinetti Balance Assessment Tool) was suggested [13]. Therefore, the result of sample size calculation of the current study suggests that a sample size of 15 older adult participants (effect size = 1.1, $\alpha = 0.05$, power = 0.8) per group was required for sufficient power to detect the difference in terms of the improvement of balance ability between participants in the Ai Chi exercise and control groups. Conservatively, we plan to recruit 30% more participants to account for potential dropouts. As a result, 20 older adult participants per group and total participants of 40 will be recruited for the current study.

2.3. Outcomes

The feasibility and preliminary efficacy of the Ai Chi Aquatic Exercise Group and the Land-based Exercise Group will be assessed. Feasibility of the Ai Chi Aquatic Exercise Group will be examined quantitatively and qualitatively in the context of recruitment, adherence, retention, feedback, subjective exercise experiences and satisfaction of the participants. Recruitment will be assessed by the proportion of potential applicants that remain interested after randomization and being informed about the procedures of the whole study. Adherence will be measured by means of attendance to exercise sessions and compliance to the exercise styles of the participants in different groups. Retention will be determined by the attrition rate as discontinuation of the intervention by different exercise groups. A

structural assessment form will be used to record all participants' feedback. All Participants' subjective exercise experiences and satisfaction will be measured by a Subjective Exercise Experience Scale and a Participant Satisfaction Questionnaire, respectively.

Additionally, preliminary efficacy of the two exercise groups will be identified through examining the mitigation of various physical and psychological fall risk factors. A series of assessment will be completed to assess various physical and psychological fall risk factors of the participants in all two assessment sessions (T0 & T1). Physical fall risk factors assessed will include flexibility, muscle strength, and clinical gait and balance. Flexibility will be measured by the 'sit-and-reach' test [17]. Longer reaching distance represents better flexibility. Muscle strength of the major lower limb's muscles groups (e.g., knee & hip extensors) will be measured using the well-recognized Lafayette manual muscle testing system (e.g., Model 01185). Higher score represents better muscle strength. Clinical gait and balance assessment will be done by the Tinetti Balance Assessment Tool [15], Timed 'Up & Go' Tests (TUG) [18] and the Berg Balance Scale (BBS) [19]. The Tinetti Balance Assessment tool assesses gait (12 points) and balance (16 points) components (i.e., total score of 28 points) at the same time [16]. Lower score represents higher risk of falling. The TUG requires participants to stand up from a chair, walk 3 m at a comfortable speed, turn around, walk back for 3 m, turn around and then sit down on the chair. Longer time to complete the TUG represents higher risk of falling. The BBS consists of 14-item tasks (daily activities) in which participants need to maintain balance in performing the tasks and each item can be scored 0 to 4 marks giving a maximum total score of 56. Lower score represents poorer balance ability and therefore higher risk of falling. Psychological fall risk factors assessment will include preliminary cognitive function, anxiety level, level of depression and fear of falling. The Chinese version Mini-Mental State Examination (MMSE-C) [16] will be used to evaluate the preliminary cognitive function at the beginning of training as a screening tool. Anxiety level will be measured by the valid Chinese version of the State-Trait Anxiety Inventory [20]. Higher score represents higher anxiety level and therefore higher risk of falling. Level of Depression will be assessed by the Chinese version 4-item Geriatric Depression Scale (GDS-4C) [21]. The GDS-4C is a reliable and valid psychological measurement of depression in older adults and was demonstrated to be an excellent alternative of the 15-item and 30-item version of the Geriatric Depression Scale with good sensitivity (80–76%) and specificity (85–81%) using the cut-off point of 2 or more [21]. Higher score represents higher level of depression and therefore higher risk of falling. The Chinese version of the Falls Efficacy Scale International (FES-I (Ch)) [22] will be implemented to assess the fear of falling. Higher score represents higher fear of falling and therefore higher risk of falling.

2.4. Intervention

Participants in both Ai Chi Aquatic Exercise Group and Land-based Exercise Group will be invited to participate in their assigned groups sixteen exercise sessions (about 60 mins each) twice per week for a total of 9 weeks (Fig. 1). All training sessions will be conducted by two experienced certified Ai Chi trainers (Ai Chi Aquatic Exercise Group) or two experienced certified fitness instructors (Land-based Exercise Group) under the supervision of a Registered Physical Therapist in Hong Kong who is required to have the professional qualifications of both Ai Chi and exercise specialists. During the group exercise session, participants will practice either a nineteen-style Ai Chi aquatic exercise program or a nineteen-style land-based exercise program, depending on their randomly assigned group. The two exercise programs were developed by a group of qualified Ai Chi and rehabilitation experts in Hong Kong based on the Ai Chi styles that was originated from Mr. Jun Kanno or the Ottago Exercise program to prevent falls in older adults, respectively. In the Ai Chi Aquatic Exercise Group, participants practised the 19-style aquatic exercise program of 1, Contemplating, 2,

Floating, 3, Uplifting, 4, Enclosing, 5, Folding, 6, Soothing, 7, Gathering, 8, Freeing, 9, Shifting, 10, Accepting, 11, Accepting with grace, 12, Rounding, 13, Balancing, 14, Flowing, 15, Reflecting, 16, Suspending, 17, Encircling, 18, Surrounding, and 19, Nurturing. In the Land-based Exercise Group, participants practised the 19-style of Ottago exercise program of 1, Trunk Movements, 2, Back Extension, 3, Ankle Movements, 4, Front Knee Strengthening Exercise, 5, Stand to Sit (No Hands), 6, Back Knee Strengthening Exercise, 7, Side Hip Strengthening Exercise, 8, Calf Raises (No Support), 9, Toe Raises (No Support), 10, Knee Bends (No Support), 11, Backwards Walking (No support), 12, Walking and Turning Around, 13, Sideways Walking, 14, Heel Toe Standing (No Support), 15, Heel Toe Walking (No Support), 16, One Leg Standing (No Support), 17, Heel Walking (No Support), 18, Toe Walking (No Support), and 19, Heel Toe Walking Backwards.

2.5. Data processing and analysis

Intention-to-treat (ITT) analysis will be adopted. For the feasibility, descriptive statistics will be used to compare the recruitment, adherence, retention and feedback of the participants in the Ai Chi Aquatic Exercise Group and the Land-based Exercise Group. Independent t-tests will be employed to compare the scores of the Subjective Exercise Experience Scale and Participant Satisfaction Questionnaire between the Ai Chi Aquatic Exercise Group and the Land-based Exercise Group.

For the preliminary efficacy, Analysis of Variance (ANOVA) with repeated measures and additional statistical tests, will be utilized to investigate the within group differences (i.e., T0 and T1) and between group differences (i.e., Ai Chi Aquatic Exercise Group & Land-based Exercise Group) in all physical and psychological fall risk factors (i.e., flexibility, muscle strength, scores of the clinical gait and balance assessments, anxiety level, level of depression and fear of falling). We could also determine the Minimal Clinically Important Differences (MCID) of the primary outcome measures (e.g., BBS) in order to help us know more about whether the intervention can lead to reducing fall risk factors.

3. Discussion

It is not uncommon among older adults, both in Hong Kong and worldwide, to suffer from falls and even fall-related injuries that required hospitalization. Given the progressively accumulating evidence to support the role of aquatic exercises for fall prevention in older adults, an expertly-designed Ai Chi aquatic exercise program could provide an alternative exercise choice for older adults who are at risk of falling. Therefore, our current study attempts to examine the feasibility and preliminary efficacy of the Ai Chi aquatic exercise program that was developed by a group of Ai Chi and rehabilitation experts in Hong Kong. If the results can provide evidence that the Ai Chi aquatic exercise program is feasible and can significantly mitigate fall risk factors in Hong Kong's older adults with moderate to high risk of falling compared to the land-based Exercise program, the structured Ai Chi aquatic exercise program could be considered to be incorporated into the standard fall prevention routine program of the Hong Kong's and even worldwide health care system.

The strength of our study is that it represents the first attempt to conduct a randomized controlled trial to examine the feasibility and preliminary efficacy of the expertly-designed Ai Chi aquatic exercise program that was developed by a group of Ai Chi and rehabilitation experts in Hong Kong. The result could provide a foundation to inform a large scale and cross-country randomized controlled trial to further investigate the effectiveness of the Ai Chi aquatic exercise program for prevention of falls in older adults in different countries. The current study may therefore with considerable global public health significance as the results could enhance our contemporary understanding of the structural Ai Chi aquatic exercise program with high practical




Fig. 1. Schematic diagram of the study design (randomized controlled trial).

implication for older adults. Based on the results of the current study, a large scale and cross-country clinical trial would be needed to further assess and confirm the effectiveness of the Ai Chi aquatic exercise program in different culture before it could be fully incorporated in the routine fall prevention program.

In conclusion, this study contributes to evaluating the effectiveness of two imperial community fall prevention exercise programs in Hong Kong. By comparing the feasibility and preliminary efficacy of the programs (i.e., the Ai Chi aquatic exercise program and the land-based exercise program), the feasibility, effectiveness and sustainability of the programs for older adults in Hong Kong could be better understood, benefiting future design of the fall prevention protocol. Consequently, this research will provide a contemporary contribution to the global evidence-based practice of fall prevention in older adults by Physical Therapists. Future follow-up study could consider evaluating the long term adherence of the Ai Chi aquatic exercise program.

Ethics approval

Ethical approval was obtained from the Institutional Review Board of the University of Hong Kong/Hospital Authority Hong Kong West Cluster (IRB Reference Number: UW 18-065). The informed consent will be obtained for eligible participants. At the first meeting with all participants, the study protocol and informed consent form will be described in detail, where all participants will be offered the opportunity to review and ask questions. An incentive of \$50 (Hong Kong Dollars) will be paid to each participant as reimbursement for costs that are associated with travelling in each attended session.

Funding support

This research is funded by the Health and Medical Research Fund, Research Fund Secretariat, Food and Health Bureau, The Government of the Hong Kong Special Administrative Region of the People's Republic of China (Grant Number: 15162131).

Clinical trial registration

The trial was registered in the HKU Clinical Trial Registry (HKUCTR-2528).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.cct.2019.100376>.

References

- [1] R.A. Marshall, J.S. Stevens, S. Velinder, T.J. van der Griend, P. Parks, *Injuries in fall-related hospitalizations in the United States, 2001–2003*, *J. Trauma Acute Care Surg.* 71 (12) (2011) 258–259.
- [2] L.W. Chu, I. Chi, A.Y. Chan, *Incidence and predictors of falls in the Chinese elderly*, *Ann. Acad. Med. Singapore* 34 (2005) 50–55.
- [3] S.M.S. Dunn, J.C. Chan, A.S.W. Wong, S.J. Lord, *Falls incidence, risk factors, and consequences in Chinese older people: a systematic review*, *J. Am. Geriatr. Soc.* 59 (2011) 530–540.
- [4] The Hong Kong Hospital Authority, *Hospital Authority Statistical Report 2014–2017*, Hong Kong Special Administrative Region, People's Republic of China, 2018.
- [5] J. Wu, Y.F. Chan, *Aging in Hong Kong: the institutional population*, *J. Am. Med. Dir. Assoc.* 10 (7) (2003) 478–483.
- [6] J.D. Gillespie, M.C. Robertson, W.J. Gillespie, S.F. Lamb, S. Gates, R.G. Cumming, R.J. Rowe, *Interventions for preventing falls in older people living in the community*, *Cochrane Database Syst. Rev.* 2 (2009) 7145.
- [7] C. Stratigopoulos, J.C. Whitley, S.H. Lord, R.J. Herbert, R.W. Cummings, J.G. Clark, *Effective exercise for the prevention of falls: a systematic review and meta-analysis*, *J. Am. Geriatr. Soc.* 56 (2008) 2234–2243.
- [8] J.L. Riggs, D.W. Browner, W.J. Hayes, R.B. Marcus, E.F. Orloff, L.J. Melton, *Differences in bone mineral density of the appendicular and axial skeleton with aging: relationship to age, sex, and weight*, *J. Clin. Invest.* 67 (1981) 238–245.
- [9] J.A. Hale, D. Waters, P. Robinson, *A randomized controlled trial to investigate the effect of water-based exercise to improve fall risk and physical function in older adults with lower-extremity osteoarthritis*, *Arch. Phys. Med. Rehabil.* 93 (2012) 27–34.
- [10] S.E. Lord, B. Mithras, R. St George, *The effects of water exercise on physical functioning in older people*, *Australas. J. Aging* 25 (2006) 36–41.
- [11] T. Teeratrakul, A. Teitl, K. Dible, A. Zafarideh, S. Wells, *The effects of a twenty-four-week aquatic training Programme on muscular strength performance in healthy elderly women*, *J. Strength Cond. Res.* 20 (2006) 851–858.
- [12] K. Sasa, J. Kozuka, *Ai Chi* Paracenter, *Aquat. Ther.* 3 (2002) 19–23.
- [13] T. Teitoku, L. Wong, J. Lambeck, P. Nieu, *The influence of Ai Chi on balance and fall risk in older adults: a randomized clinical trial*, *Physiotherapy* 97 (2017) 1011–1019.

- [14] S.I. Morales-Gonzalez, L. Viqueiro-Torres, J.D. Martinez-Alba, Whole-body Tai Chi chiao-chun benefits in musculoskeletal disease: current evidence, *J. Bonec. Rehabil.* 11 (3) (2015) 120.
- [15] H.T.K. Chiu, H.C. Lee, W.S. Chang, R.K. Kuo, Reliability and validity of the Cantonese version of Tai Chi-mental awareness index: a preliminary study, *J. Hong Kong Coll. Physiother.* 4 (1998) 25–28.
- [16] M.E. Tietini, Performance-Oriented assessment of mobility problems in elderly patients, *J. Am. Geriatr. Soc.* 34 (1986) 119–120.
- [17] J. Clark, W. Chuang, R. Adams, W.W.K. Wong, D. East, B. Whinnell, Tests for fitness in older adults: MIMEDS fitness task force, *J. Phys. Educ. Recreat. Dance* 69 (1999) 50–71.
- [18] D. Pedrinelli, L. Richardson, The Tamed “Up & Go”: a test of basic functional mobility for frail elderly persons, *J. Am. Geriatr. Soc.* 39 (1991) 142–146.
- [19] E. Berg, S. Wood-Campness, J.I. Williams, U. Gostono, Measuring balance in the elderly: preliminary development of an instrument, *Physiother. Can.* 43 (1990) 304.
- [20] D.T. Shah, The factorial structure of the Chinese version of the State-Trait Anxiety Inventory: a confirmatory factor analysis, *Behav. Psychol. Mon.* 61 (1991) 685–697.
- [21] S.T. Cheung, A.C.W. Chan, Cross-race performance of long and short forms of the Geriatric Depression Scale in mildly depressed Chinese, *Int. J. Geriatr. Psychiatry* 23 (2008) 1121–1137.
- [22] S.M.S. Egan, W.P.N. Tsang, J.C.T. Chan, S.R. Lord, Development and validation of a Chinese version of the Tai Chi-ay male instrument, *Arch. Gerontol. Geriatr.* 58 (2013) 159–174.

Effect of Tai Chi Exercises on the Balance, Functional Gait, and Flexibility of Elderly Filipino Males

Sheng Chen^{1,a}

¹College of Physical Education, Chengdu University, Sichuan Chengdu, China

^a787345913@gmail.com

Keywords: Tai Chi

Abstract: The objective of this study is to evaluate the effect of a 16-week Tai Chi Chuan exercise program on the balance, functional gait and sit-and-reach flexibility among selected elderly Filipino males. This research utilized a one-group pre-test and post-test quasi-experimental design. Forty healthy male Filipino respondents aged 55-60 years, with no prior experience or knowledge about Tai Chi Chuan, were recruited from an urban community in Pasay City, Philippines. All of the respondents participated in a 16-week exercise program that was held three times a week every morning. Balance, functional gait and sit-and-reach flexibility were measured at the beginning and the end of the Tai Chi Chuan exercise intervention. This study showed that Tai Chi offers potential benefits to the elderly males in terms of balance and other physical functions. Improvements were evident in functional balance, gait and flexibility. The study proves that Tai Chi should be an alternative method for elderly males to slow down the normal decline in balance, gait performance and physical flexibility.

1. Introduction

Tai Chi Chuan (Taijiquan) is a healing martial art that combines martial arts movements with energy circulation known as *qi* or *chi*, breathing, and stretching Techniques [1]. Originally developed for self-defense, tai chi has evolved into a graceful form of exercises that's now used for stress reduction and a variety of other health conditions (Mayo Clinic, 2015). As a proven life-prolonging, healing, and rejuvenating exercise, Tai Chi have been found to be ideal for older people (Liang and Shu, 2005; The Telegraph, 2012). Researchers found that older people who regularly performed the traditional Chinese mind and body technique were less likely to suffer health problems and were physically stronger (Yeh, Wang, Wayne, Phillips, 2008). Tai Chi is ideal for the elderly as it is a low-impact, slow-motion exercise in which a person moves while breathing deeply and naturally, thus, focusing one's attention on bodily sensations, which is similar to some kinds of meditation (Harvard Health, 2015). In addition, Tai Chi has been proven to be beneficial in lowering certain risks that older people are most prone to. Such is the case for their vulnerability to falls which has been an important cause of morbidity among the elderly (Santos, da Silva, de Pinho, Gautério, Pelzer, and da Silveira, 2012). Santos et al. (2012) added that one of the most important physiological factors for falls is impaired balance. Many factors contribute to poor balance, including reduced strength, flexibility, and sensorimotor coordination as well as delayed information processing (Yu and Yang, 2012).

There are several cross-sectional and experimental studies proving how Tai Chi benefits the elderly by improving their balance and physical fitness. In 2012, Yu and Yang found that Tai Chi improved the balance and other physical functions such as flexibility and reaction time of elderly male subjects. The positive effect of Tai Chi on balance were further supported by the study of Konig, Galarza, Goulart, Lanferdini, Tiggeman, and Dias (2014). Their study saw significant improvement in balance through different evaluation methods after three and six months of Tai Chi intervention among elderly women. In an even earlier study, Li, Hamer, Fisher, and McAuley (2004) found that older people who practiced Tai Chi performed significantly better compared to non-practitioners. In addition, while both groups in their study exhibited deterioration in functional

balance measures during post-intervention follow-up, the Tai Chi group showed a significantly slower rate. Other studies proved the benefits of Tai Chi among elderly people by significantly improving different health indices such as blood pressure (Yeh, et al., 2008) and cognition (Chang, Tsai, Beck, Hagen, Huff, Anand, Roberson, Rosengren, and Beuscher, 2011). The findings of these studies support the health benefits of Tai Chi as a form of exercise among elderly men and women. While most studies have focused on evaluating Tai Chi through balance tests, this study aimed to further support Tai Chi's positive effect on other factors such as flexibility and functional gait that may affect balance.

The objective of this study was to evaluate the effect of a 16-week Tai Chi Chuan exercise program on the balance, functional gait, and sit-and-reach flexibility among selected elderly Filipino males. In due course, it is hypothesized that the Tai Chi Chuan intervention conducted three times a week for one hour per session in a span of 16 weeks was sufficient to produce positive changes among the target respondents' balance, functional gait, and flexibility.

2. Methods

2.1. Research Design and Sampling

This research utilized a one-group pre-test and post-test quasi-experimental design. Forty male Filipino respondents aged 55-60 years, with no prior experience or knowledge about Tai Chi Chuan, were recruited from an urban community in Pasay City, Philippines. The exclusion criteria included the presence of health problems such as cognitive impairments, symptomatic cardiovascular diseases at moderate exertion levels, poorly controlled hypertension or symptomatic orthostatic hypotension, other neurological disorders, peripheral neuropathy of the lower extremities, crippling arthritis, and metastatic cancers. In addition, the respondents were not taking any medications that could affect their balance. All of the respondents were asked to refrain from changing their daily lifestyle routine. Informed consents were obtained from the respondents after thoroughly explaining the objectives and procedures of the study, including the benefits and risks of participation.

2.2. Exercise Intervention

All of the respondents participated in a 16-week exercise program that was held three times a week every morning. Each Tai Chi Chuan exercise session lasted 60 minutes. A certified Tai Chi Chuan instructor with one assistant facilitated each session in which the 24-posture simplified Tai Chi Chuan was utilized. In each session, the first 10 minutes were dedicated for warm-up exercises, the next 40 minutes were for the main Tai Chi Chuan exercise, and the concluding 10 minutes were reserved for cool-down exercises. Correct body positioning, joint angles, and form-to-form transitions were constantly monitored during the sessions.

2.3. Outcome Measures

Balance, functional gait, and sit-and-reach flexibility were measured at the beginning and the end of the Tai Chi Chuan exercise intervention. While functional gait and flexibility have been identified as contributing factors to balance, all three are important factors that have been associated with risk of falls among the elderly (Yu and Yang, 2012; Kouig, et al., 2014).

2.4. Balance

Functional balance among the elderly respondents was evaluated using the Berg Balance Scale (BBS). It is based on 14 items common to everyday life. The maximum score that can be achieved in this scale is 56 and each item has an ordinal scale consisting of five options ranging from 0 to 4 points according to the level of difficulty (the higher the score, the better the balance). The test is simple, easy to administer, and safe for the evaluation of elderly subjects. (Berg, Wood-Dauphinee, Williams, and Gayton, 1989; Miyamoto, Lombardi, Berg, Natour, and Ramos, 2004)

2.5. Functional Gait

The 'timed up and go' test (TUG) is a simple, quick and widely used clinical performance-based measure of lower extremity function, mobility, and fall risk (Herman, Giladi, and Hausdorff, 2011). TUG is one of the more commonly used clinical tools for quantifying gait, dynamic balance abilities

and fall risk. In this study, TUG was mainly used to measure functional gait. It is a simple test in which the elderly respondents were asked to stand up from a chair (resting position), walk a distance of three meters and then return to the chair and sit again (with their backs on the backrest). The respondents were instructed to perform the task safely and as quickly as possible and their performances were analyzed by counting the time they needed to complete the test (Kouig, et al., 2014). Before each test, a familiarization protocol was conducted so as to ensure safety and accurate results.

2.6. Sit-and-Reach Flexibility

Sit-and-reach flexibility was measured using a sit-and-reach apparatus. The evaluation protocol was patterned with the study of Yu and Yang (2012). All of the elderly respondents were asked to sit on the floor with their legs stretched out forward and their shoes removed. Both knees were locked and pressed flat to the floor (the tester assisted by holding them down). With the palms facing downward, the subjects reached forward along the measuring line as far as possible and held that position for one or two seconds while the distance was recorded in centimeters. The subjects were tested twice, and the better results were used.

2.7. Statistical Analysis

Descriptive Statistics was used (weighted mean) for the elderly respondents for both the pre and post-test measures. Inferential statistics, specifically, paired t-test was used in finding out the effect of the Tai Chi Chuan exercise intervention by identifying if there was a significant difference in the outcome measures between the pre-test and the post-test. The level of significance was set at $p < 0.05$.

3. Results

Figure 1 presents the mean of the outcome measure pertaining to balance pre- and post-intervention. After the 16-week Tai Chi Chuan exercise intervention, functional balance significantly improved ($p = 0.02$).




Figure 1 Comparison of functional balance before and after the 16-week Tai Chi Chuan intervention

Figure 2 shows the mean of the outcome measure pertaining to functional gait pre- and post-intervention. Functional gait, which was tested using the Timed Up & Go test, is statistically better after the respondents underwent the Tai Chi intervention ($p = 0.00$).




Figure 2 Comparison of functional gait before and after the 16-week Tai Chi Chuan intervention

Figure 3 illustrates the mean of the outcome measure pertaining to flexibility pre- and post-intervention. After the 16-week Tai Chi intervention, the sit-and-reach flexibility of the respondents significantly improved ($p = 0.02$).




Figure 3 Comparison of measure of flexibility before and after the 16-week Tai Chi Chuan intervention

4. Discussion

This study shows that Tai Chi offers potential benefits to the elderly in terms of balance and other physical functions. Improvements were evident in functional balance, gait, and flexibility. Tai Chi Chuan or Tai Chi is a widely practiced Chinese martial art. It consists of a series of postures combined in slow, smooth, graceful movements and is considered a low intensity exercise that is claimed to develop balance, coordination, and help maintain strength and emotional health (Wu, 2002).

Balance is a required component in the execution of postural control. The capacity to maintain balance decreases with age which results in an increased risk of falls and fractures among the elderly (Nicken, 2010). These findings suggest that Tai Chi exercise may positively influence balance

abilities among older males during the course of 16 weeks of training. Furthermore, flexibility is an important factor in maintaining balance. One possible explanation for this finding is that Tai Chi is a mind-body practice that combines meditation with slow, gentle, graceful movements. It is considered a complex, multi-component intervention that integrates physical, psychosocial, emotional, spiritual, and behavioural elements (Wayne, et al., 2011) and features constant swinging, shifting, and turning in all directions, including left, right, forward, and backward. This activity requires a high degree of concentration and coordination between mind and body and among the different body parts. The motor system, the nervous system and the proprioceptive systems are all mobilized during Tai Chi exercise.

The results of this study proved that Tai Chi could improve the ability to stay balanced and increase muscle strength and flexibility. Moreover, it was also found out that Tai Chi is a form of exercise that can improve the flexibility of the body and mind, as well as adjust breathing, and can increase the cooperation among the hands, eyes and whole body with the mind. Tai Chi can boost the control of the mind on the body and improve the reaction capacity of the respondents. This study shows that the respondents who were subjected to Tai Chi for 16 weeks possessed better trunk and hamstring flexibility than they experienced during their previous sedentary lifestyles. This finding is confirmed by the findings of Lan et al. (2012), who reported significantly superior performance among older Tai Chi practitioners with more than 10 years of experience than their sedentary counterparts with respect to hip joint flexibility, as measured by their stand and reach test scores. Lan et al., et al (2015) found out in their study that the maintenance and development of levels of flexibility closely related to balance are important components of a general health enhancement program during the aging process.

Similarly, Chyu, et al. (2010) also found out in their study that Tai Chi exercises incorporate whole body exercises and some isotonic exercises. Performing Tai Chi exercises can help build muscles on the back and the lower limbs, which can subsequently increase muscle strength. On the other hand, Taylor, et al. (2012) discussed about the steps of Tai Chi which focus on the exchange between deficiency and excess. The so-called 'deficiency and excess' (Xu Shi in Chinese) is the proportion of the body weight excreted on the feet, the foot that withstands more body weight is referred to as 'excess,' and the foot with less body weight is referred to as 'deficiency.' 'The exchange between deficiency and excess' actually refers to the change of body weight between each foot. This theory fully emphasizes the capacity to control the distribution of the body weight and adjust the gestures. With the guidance of this theory and cooperation of a series of movements made while standing with a single leg, standing with two legs, or standing with the other leg, the capacity of the elderly to stay balanced and adjust their steps is improved.

The results indicate that, after the 16-week intervention period, functional gait, which was tested using the Timed Up & Go test, is statistically better after the respondents underwent the Tai Chi intervention ($p = 0.00$). Moreover, the sit-and-reach flexibility of the respondents also significantly improved ($p = 0.02$). It is worth mentioning that the value of the gait speed of the respondents at the baseline in the present study were consistent with those reported in a previous study (Maki, 2007). Wayne et al. (2015) reported that the gait speeds for healthy adults have been shown to range from 1.05 to 1.43 m/s. Reductions in gait speed, stride length, ranges of motion, and force momentum in hip and knee joints are common in older adults [34,35,36]. In particular, the reduction in freely selected gait speed is an indicator of risk of falls [37]. Previous studies have shown that the kinematic parameters of the knee and ankle joints have a positive relationship with gait speed (Kirkwood, et al., 2007). In particular, Kwon, et al. (2015) found that between the pre-swing and the mid-swing, the peak values of the flexion and the external rotation of the knee joint and the peak values of the plantar-flexion of the ankle joint significantly increased with the increase of gait speed.

Meanwhile, a large number of studies have shown that strength training programs have a beneficial impact on gait efficacy (Lamoureux et al., 2013), including slowing down the decline

in gait function due to age. However, with regard to hip flexibility, a few studies have shown that the stretching training program is better than the strength training program to retard the normal, age-related contraction of the muscle groups of the pelvis and hip, like the gluteal and rotator (Christiansen, et al., 2008). The results of the study indicate that the 16-week helped in improving the balance, functional gait, and sit-and-reach flexibility among selected elderly Filipino males. This probably is associated with the "Loosen Waist and Hips (Song Yao Song Kua)" guideline of Tai Chi. From a martial arts standpoint, when the waist and hips are loose, the power generated by the lower limbs can easily be transmitted to the arms. To adhere to this principle, the upper body must be upright and the stance must be comfortable, which may be effective in improving the flexibility of the hip and ankle. Lastly, the Tai Chi pattern includes many movements, which require people to go down on their knees and their waists; this is good for many joints because it increases the movement of the joints and improves flexibility (Lelard, et al. 2010).

However, psychological gains from Tai Chi have been demonstrated only in the elderly, and there is a need therefore to establish benefits in younger age cohorts for whom balance is seldom problematical. In general, there is a paucity of research on the benefits of Tai Chi for the non-elderly population, possibly because of the unwarranted perception that Tai Chi is an exercise appropriate only for the elderly.

5. Conclusion

In conclusion, Tai chi is effective in improving balance, functional gait, and sit-and-reach flexibility among selected elderly Filipino males. Therefore, the Tai Chi Chuan intervention conducted three times a week for one hour per session in a span of 16 weeks was sufficient to produce positive changes among the target respondents' balance, functional gait, and flexibility.

Given the results of the study, it should be thought of as an alternative method for older adults to slow down the normal decline in gait performance and physical flexibility. Future studies should run a similar randomized controlled trial with a large sample size to confirm the beneficial effects of Tai chi on gait parameters and musculoskeletal flexibility in different populations such as healthy female older adults, and those older adults with high risk for falls and those inflicted with disorders like Parkinson's disease and peripheral neuropathy.

References

- [1] Liang, S., Wu, W.C. (2005). *Simplified Tai Chi Chuan: 24 Postures with Applications and Standard 48 Postures*. BERTRAMS.
- [2] Nicken, H. (2010). *Intrinsic factors in falling among the elderly*. Arch Intern Med, 145 (1985), pp. 1089-1093.
- [3] Berg, K.O., Wood, D., Williams, J.L., Gaytaon, D. (2010). Measuring balance in the elderly: preliminary development of an instrument. *Physiother Can* 1989;41(6):304-11.
- [4] Chang, J., Beck, C., Hagen, J., Huff, Debbie, et al (2011). The Effect of Tai Chi on Cognition in Elders with Cognitive Impairment. *MedsurgNurs*. 2011 Mar-Apr; 20(2): 63-70. Pitman, NJ :Jannetti Publications, Inc.
- [5] Christiansen, C.I.. (2008) The effects of hip and ankle stretching on gait function of older people *Arch. Phys. Med. Rehabil.* 89, 1421-1428.
- [6] Chyu, M.C., James, C.R., Sawyer, S.F., et al (2010). Effects of tai chi exercise on posturography, gait, physical function and quality of life in postmenopausal women with osteopaenia: a randomized clinical study. *Clin Rehabil*, 24 (2010), pp. 1080-1090.
- [7] Ding, H., Hui-Xin, Y. (2012). The effect of Tai Chi intervention on balance in older males. *Journal of Sport and Health Science Volume 1, Issue 1, May 2012, Pages 57-60*.
- [8] Herman, T., Hausdorff, J. (2011). Properties of the "Timed Up and Go" Test: More than Meets the Eye. *Gerontology*. 2011 Apr; 57(3): 203-210.

Effects of Tai Chi Chuan and Brisk Walking Exercise on Balance Ability in Elderly Women: A Randomized Controlled Trial

Wei Sun

Shanghai University of Sport
 and Shandong Institute
 of Sport Science

Xiujie Ma

Shanghai University of Sport
 and Handan University

Lin Wang

Shanghai University of Sport

Cui Zhang

Shanghai University of Sport
 and Shandong Institute
 of Sport Science

Qipeng Song and Houxin Gu

Shandong Institute
 of Sport Science

Dewei Mao

Shanghai University of Sport
 and Shandong Sport University

This study aims to investigate the effects of Tai Chi Chuan (TCC) and brisk walking (BW) on balance and training duration for the two exercises to significantly improve balance. A total of 48 elderly women were randomly divided into three groups. The TCC and BW groups completed a 60-min intervention training program with five sessions weekly for 16 weeks. Single-leg standing balance was tested every 4 weeks. Results showed that all the variables with eyes open improved on the eighth week ($p < .05$) in the TCC group and on the 12th week ($p < .01$) in the BW group. All variables with eyes closed improved on the 12th week ($p < .01$) in the TCC group and on the 16th week ($p < .05$) in the BW group. The results showed that 12 and 16 weeks of TCC and BW, respectively were essential to improve balance with eyes closed among the women aged 60–70 years.

Keywords: aging, balance control, fall, intervention

Sun, Ma, Wang, Zhang, and Mao are with the Shanghai University of Sport, Shanghai, China. Sun, Zhang, Song, and Gu are with the Shandong Institute of Sport Science, Jinan, Shandong, China. Ma is also with the Handan University, Handan, China. Mao is also with the Shandong Sport University, Jinan, Shandong, China. Address author correspondence to Dewei Mao at m_d_wei@sina.com.

Approximately one of three people aged 65 years and older fall at least once a year, and half of them experience multiple falls (Davis et al., 2010). Falls cause severe injuries in the elderly, especially postmenopausal women with osteoporosis; such injuries include soft tissue injury, fracture, and brain trauma, which can result in death (Konak, Kibar, & Ergin, 2016). Decreased balance with single-leg stance (SLS) was found to be a significant predictor of falls (Enderlin et al., 2015). Falls during SLS, such as stepping over obstacles and climbing stairs, were reported to account for approximately 50% of falls in this population (Novak, Komisar, Maki, & Fernie, 2016). Balance ability is related to mobility, vision, muscle strength, and proprioception (Enderlin et al., 2015); however, the elderly experience a significant decline in organ function, such as poor eyesight, ankle and knee joint proprioception, and muscle strength. This decline in organ function that comes with aging might reduce the ability of older adults to balance with SLS and could increase their risk of falling (Huang & Liu, 2015).

Balance ability should be improved to prevent falls and injuries in the elderly (Kathiresan, Jali, Aliqah, Aznie, & Fidieyana, 2010). Moderate aerobic exercise positively affects the balance ability of older adults (Dixit, Maiya, Shastry, & Guddattu, 2016). Tai Chi Chuan (TCC), a traditional Chinese exercise form known for its slow and graceful movements, has become one of the most popular exercises in China (Chan et al., 2016). Some evidence indicates that short- and long-term TCC exercises can improve the body balance of adults and patients (Kim, Kim, & Lee, 2015; Li et al., 2012; Scianni, 2015; Song et al., 2017; Tsang, 2013; Zhou et al., 2015). Furthermore, regular TCC exercise can enhance joint kinesthesia (Chang et al., 2016; Zhang, Sun, Yu, Song, & Mao, 2014), lower extremity muscle strength (Xu, Hong, & Li, 2008), and neuromuscular reaction in elderly women (Sun et al., 2016; Wang, Xu, & Li, 2017). Brisk walking (BW), with its speed being higher than that of normal walking, is one of the most common moderate aerobic exercise forms with 40–59% maximal oxygen intake (Murtagh, Borcham, & Murphy, 2002). BW is recommended to all age groups as an exercise that requires minimal special skills and facilities. Although a number of studies have indicated that BW can enhance static and dynamic balance that reduces fall risk among elderly adults (Okubo et al., 2015; Paillard, Lafont, Costes-Salon, Rivière, & Dupui, 2004), other studies have yielded inconsistent results about the positive effects of a 10-week BW intervention on static balance ability (Gába et al., 2016).

Regular TCC exercise could improve balance control, but the essential training duration for improvements remains unclear. Moreover, the few studies on the effects of BW on balance ability in the elderly have shown contradictory results. The present study aims to assess the training effects of TCC and BW exercises on balance ability and to identify the ideal training duration necessary for the two exercises to significantly improve balance. We hypothesize that TCC exercise is more efficient than BW exercise in improving body balance in SLS under two visual conditions.

Methods

Study Design

A randomized controlled trial was designed to compare the effects of TCC and BW exercises on balance with SLS during a 16-week training program. Both TCC and

(Ahead of Print)

BW groups participated in a 60-min moderate-intensity intervention exercise for at least five sessions weekly for 16 weeks. Balance ability was tested at the baseline and on the 4th, 8th, 12th, and 16th week.

Recruitment of Study Participants

A total of 48 elderly women (aged 60–70 years) were recruited by means of advertisements in newspapers and leaflets and through community advocacies in Jinan, Shandong Province, China. The exclusion criteria are as follows: any regular exercise experience; falling history; and cardiovascular, neurological, and musculoskeletal diseases. The application for human ethics and informed consent forms was approved by the medical ethics committee of Shandong Institute of Sport Science.

Randomization

After baseline assessments, all participants were randomly divided into TCC, BW, and control groups with the use of permuted block randomization at a ratio of 1:1:1. The randomized list, sealed in an envelope, was provided to the researcher before implementing the intervention.

Intervention

In the first 3 weeks, the participants were individually instructed about the 24-form TCC exercise by a qualified TCC master. Each session included a 10-min warm-up, a 20-min training on new movement forms, a 20-min review of learned movements, and a 10-min cooldown. The participants then practiced under the supervision of a master for the next 13 weeks. Each session included a 10-min warm-up, a 40-min TCC exercise, and a 10-min cooldown.

BW is defined as walking at a speed of 1.79 m/s in an indoor track (Murtagh et al., 2002). The participants in the BW group walked at a speed perceived to cause breathing to accelerate and the body to warm-up and sweat (Gába et al., 2016). The pace and speed on a pedestrian road were set by the professional instructor. The walking duration increased from 10 to 40 min over the first 3 weeks. It was then maintained at 40 min for the next 13 weeks. Each session consisted of a 10-min warm-up, a 40-min BW exercise, and a 10-min cooldown.

The control group was organized to watch TV programs, read newspapers, or attend health education lectures with the same schedule as the other two groups. However, they were prohibited from engaging in any regular exercise and allowed to maintain their own dietary habits.

Measurements

The measurements in the present study consisted of foot pressure plate and SLS time measurements. SLS tests, with an internal consistency reliability ranging from 0.95 to 0.99 (Franchignoni, Tesio, Martino, & Ricupero, 1998), have been extensively used to assess balance and exercise among the elderly. Both measurements were completed in a quiet testing room in the morning.

(Ahead of Print)

Foot Pressure Plate Measurement

Balance tests were performed with a foot pressure plate (RSscan for 2D balance with a 0.5-m system, Belgium; Hijmans, Zijlstra, Geertzen, Hof, & Postema, 2009). Each participant was asked to stand barefoot with the dominant leg in a comfortable self-chosen stance facing forward on a plate; the dominant leg is described as the preferred leg for kicking a football (Gribble, Tucker, & White, 2006), as motionless as possible. The other leg was fixed at 90° with the hip and knee joints flexed and both arms relaxed at the sides of the body. Two standing conditions were randomly tested: in one condition, the participants were asked to stand on a single leg for 22 s with their eyes open (i.e., SLS with eyes open, SLO) and looking straight ahead at a dot on a wall 2.5 m away (Nilsson, Ageberg, Ekdahl, & Encroth, 2006); in the other condition, they performed an SLS for 12 s with their eyes closed (i.e., SLS with eyes closed, SLC; Gerbino, Griffin, & Zurakowski, 2007). The trial failed if the supporting leg moved or the nonsupporting leg touched the floor. Three successful trials for each SLO and SLC were tested after the procedures were familiarized. The time interval for breaks was 1 min between two consecutive trials. All measurement procedures were performed under the supervision of a technician.

The data were sampled at 17 Hz and low-pass filtered with a cutoff frequency of 6 Hz (Butterworth; Hijmans et al., 2009). The data from first and last seconds in the trial were not included in data analysis. On the basis of the displacement of the center of pressure (COP), all outcomes were calculated to assess postural steadiness (Prieto, Myklebust, Hoffmann, & Lovett, 1996; Figure 1). The maximal




Figure 1 — COP trajectories of SLO and SLC for a representative participant. COP = center of pressure; SLO = SLS with eyes open; SLC = SLS with eyes closed; SLS = single-leg stance; AP = anterior–posterior; ML = medial–lateral.

(Ahead of Print)

displacements of COP in the anterior–posterior (AP; D-ap) and medial–lateral (ML; D-ml) directions, as well as the total length of the COP trajectories (L_{ng}) and the 95% confidence ellipse area of COP movements (Area), were derived (Prieto et al., 1996).

SLS Time Measurement

This measurement procedure was similar to the foot pressure plate measure. The participants were asked to stand on the ground with their dominant leg in SLO and SLC and with arms positioned along the sides of the body. The participants performed SLO and SLC three times, and the set with the longest duration was selected for analysis. A 1-min interval was provided as a break between two consecutive trials.

Statistical Analysis

The software SPSS 17.0 (IBM, New York) was used for data analysis, and all the variables were presented as mean $\pm$ SD. The variables of five measurements were named week₀, week₄, week₈, week₁₂, and week₁₆ in this study. One-way analysis of variance (ANOVA) was used to compare the differences of the demographic and baseline variables across the three groups. Dunnett's T3 was chosen for post hoc multiple comparisons for nonequal homogeneity of variances. Bonferroni method was selected for equal homogeneity of variances. Two-way repeated-measures ANOVA was used to determine the effects of different groups and time durations, as well as their interaction effects, on the measurements. If any significant main and interaction effects were found, then the Bonferroni method was conducted for post hoc comparisons. The significance level was set at .05.

Results

Baseline Characteristics

A total of 50 persons were screened for eligibility; 48 were deemed qualified and then divided into three groups. A total of 36 participants (12 in the TCC group, 13 in the BW group, and 11 in the control group) completed the entire 16-week study. Furthermore, 64 attendance sessions out of 80 (80%) were required for each participant in the three groups. Twelve individuals dropped out because of health reasons ($n = 2$), low attendance rate ($n = 6$), lack of time ($n = 3$), and no reason ($n = 1$). No statistical difference in baseline characteristics was detected among the three groups (Table 1).

Balance Ability With SLS

As shown in Table 2, in SLS with eyes open, two-way repeated-measures ANOVA showed significant group effects, time effects, and Group $\times$ Time interaction effects. No significant difference in all the variables was observed among the three groups at week₀. In week₈, significant within-group differences in all the variables were observed in the TCC group. The same differences were observed in the BW group in week₁₂. The detailed post hoc statistical results are as follows:

(Ahead of Print)

Table 1 Baseline Characteristics of Participants

	TCC Group (n = 12)	BW Group (n = 13)	C Group (n = 11)	F	p
Age (years)	64.12 ± 3.21	63.26 ± 2.20	65.36 ± 4.31	0.712	.498
Weight (kg)	62.81 ± 8.37	62.00 ± 7.49	62.63 ± 7.21	0.004	.996
Height (cm)	157.56 ± 5.45	158.50 ± 4.40	156.45 ± 4.43	2.607	.089
BMI (kg/m ²)	25.12 ± 3.19	24.69 ± 2.97	26.21 ± 3.82	0.617	.546
Education (years)	6.83 ± 0.83	7.15 ± 0.80	7.27 ± 0.79	0.926	.406

Note. Data are presented as mean ± SD. TCC = Tai Chi Chuan; BW = brisk walking; C = control; BMI = body mass index.

balance ability significantly increased by 57.6% (TCC in week₈) and 35.8% (BW in week₁₂) for Time and decreased by 40.2% (TCC in week₈) and 42.5% (BW in week₈) for D-ap, by 45.2% (TCC in week₈) and 35.4% (BW in week₈) for D-ml, by 39.7% (TCC in week₈) and 28.8% (BW in week₁₂) for Lng, and by 65.3% (TCC in week₈) and 71.0% (BW in week₈) for Area relative to the baseline, respectively. No significant change from the baseline was observed in the control group. Significant differences in all the variables were observed between the TCC and control groups in week₈, whereas these differences were observed between BW and control groups in week₁₂.

As shown in Table 3, in SLS with eyes closed, two-way repeated-measures ANOVA revealed significant group effects, time effects, and Group × Time interaction effects. No significant difference in all variables was observed among the three groups at week₀. Significant within-group differences in all variables were observed in the TCC group in week₁₂ and in the BW group in week₁₆. The detailed post hoc statistical results are as follows: balance ability significantly increased by 94.3% (TCC in week₁₂) and 101.5% (BW in week₁₆) for Time and decreased by 31.7% (TCC in week₁₂) and 30.0% (BW in week₈) for D-ap, by 20.2% (TCC in week₈) and 26.8% (BW in week₁₆) for D-ml, by 36.4% (TCC in week₄) and 38.8% (BW in week₄) for Lng, and by 42.8% (TCC in week₈) and 38.1% (BW in week₁₂) for Area from the baseline, respectively. No significant change from the baseline was observed in the control group. Compared with the control group, significant between-group differences were found in Time (TCC at week₁₂ and BW at week₁₆), Lng (TCC at week₄ and BW at week₄), Area (TCC at week₁₆), and D-ap (TCC at week₁₂ and BW at week₈).

Discussion

The results of the present study, which support the hypothesis, showed that the variables of balance ability, D-ap, D-ml, Lng, Area, and Time significantly improved in the TCC and BW groups after training. The balance ability in the control group remained unchanged during the 16-week experimental period, thereby indicating no learning effects on the balance test. Thus, both TCC and BW exercises can improve balance under SLO and SLC via intervention training.

(Ahead of Print)

Table 2 Comparisons of Study Variables With Eyes Open in Single-Leg Stance Among Three Groups

	TCC Group (n = 12)	BW Group (n = 13)	C Group (n = 11)	Group		Time		Group × Time	
				F	p	F	p	F	p
Time (s)									
Week ₀	32.73 ± 16.69	40.80 ± 15.13	38.37 ± 13.73	5.21	.006	4.05	.004	1.22	.291
Week ₄	39.14 ± 21.57	41.66 ± 14.66	40.22 ± 12.15						
Week ₈	51.59 ± 15.26 ^a	44.61 ± 14.12	40.38 ± 9.58						
Week ₁₂	53.45 ± 14.23 ^{a,b}	55.44 ± 8.62 ^{a,c}	38.70 ± 12.68						
Week ₁₆	55.61 ± 10.20 ^{a,b}	55.03 ± 12.80 ^{a,c}	39.94 ± 8.05						
Lng (mm)									
Week ₀	422.57 ± 105.56	363.43 ± 104.7	427.93 ± 152.3	17.38	<.001	8.88	<.001	1.35	.219
Week ₄	357.54 ± 114.62	347.50 ± 77.92	404.56 ± 120.10						
Week ₈	257.92 ± 74.84 ^{a,b}	293.04 ± 86.61	409.0 ± 176.20						
Week ₁₂	235.62 ± 65.24 ^{a,b}	261.28 ± 98.10 ^a	386.1 ± 83.09						
Week ₁₆	217.89 ± 45.82 ^{a,b}	245.42 ± 80.48 ^{a,c}	375.9 ± 69.78						
Area (cm ²)									
Week ₀	1.45 ± 0.53	1.32 ± 0.61	1.38 ± 0.71	29.88	<.001	13.22	<.001	2.76	.007
Week ₄	0.93 ± 0.55	0.98 ± 0.52	1.34 ± 0.72						
Week ₈	0.50 ± 0.31 ^{a,b}	0.38 ± 0.25 ^{a,c}	1.32 ± 0.20						
Week ₁₂	0.56 ± 0.39 ^{a,b}	0.58 ± 0.41 ^{a,c}	1.38 ± 0.16						
Week ₁₆	0.40 ± 0.11 ^{a,b}	0.34 ± 0.25 ^{a,c}	1.29 ± 0.20						

(continued)

Table 2 (continued)

	TCC Group (n = 12)	BW Group (n = 13)	C Group (n = 11)	Group		Time		Group × Time	
				F	p	F	p	F	p
D-ap (mm)									
Week ₀	30.07 ± 4.82	28.47 ± 7.96	25.95 ± 6.71	25.54	<.001	16.58	<.001	2.25	.026
Week ₄	23.21 ± 5.13 ^b	23.02 ± 4.49	25.30 ± 8.13						
Week ₈	18.31 ± 5.72 ^{a,b}	16.42 ± 5.44 ^{a,c}	24.92 ± 7.06						
Week ₁₂	17.98 ± 5.30 ^{a,b}	17.75 ± 6.33 ^{a,c}	25.32 ± 6.00						
Week ₁₆	16.52 ± 3.62 ^{a,b}	15.60 ± 5.64 ^{a,c}	25.61 ± 6.51						
D-ml (mm)									
Week ₀	33.62 ± 11.49	31.97 ± 9.88	33.59 ± 7.32	12.08	<.001	13.78	<.001	2.95	.004
Week ₄	24.01 ± 5.23	29.30 ± 9.39	32.80 ± 4.08						
Week ₈	18.05 ± 7.21 ^{a,b}	20.22 ± 5.71 ^{a,c}	30.49 ± 5.81						
Week ₁₂	19.14 ± 4.58 ^{a,b}	21.13 ± 7.23 ^{a,c}	30.83 ± 5.92						
Week ₁₆	18.61 ± 6.49 ^{a,b}	16.46 ± 6.08 ^{a,c}	29.26 ± 3.96						

Note. Data are presented as mean ± SD. TCC = Tai Chi Chuan; BW = brisk walking; C = control; Time = the length of time with single-leg stance; Lug = the total length of the COP trajectories; Area = the area of 95% bivariate confidence ellipse, which encloses approximately 95% of the points on the COP path; D-ap = the maximal displacement of COP in the anterior-posterior direction; D-ml = the maximal displacement of COP in the medial-lateral direction; COP = center of pressure.

^aSignificant difference compared with the value at week₀ within each group. ^bSignificant difference between TCC and C groups. ^cSignificant difference between BW and C groups.

Table 3 Comparisons of Study Variables With Eyes Closed in Single-Leg Stance Among Three Groups

	TCC Group (n = 12)	BW Group (n = 13)	C Group (n = 11)	Group		Time		Group × Time	
				F	p	F	p	F	p
Time (s)									
Week ₀	16.78 ± 7.1	15.63 ± 8.3	18.26 ± 7.63	7.36	.001	11.67	<.001	2.98	.004
Week ₄	19.14 ± 7.59	18.95 ± 6.2	18.70 ± 11.66						
Week ₈	21.88 ± 8.08	23.44 ± 12.89	19.97 ± 8.36						
Week ₁₂	31.42 ± 9.23 ^{a,b}	27.08 ± 12.08	19.53 ± 9.18						
Week ₁₆	39.95 ± 11.67 ^{a,b}	31.68 ± 12.4 ^{a,c}	19.11 ± 8.19						
Lag (mm)									
Week ₀	519.51 ± 105.54	582.20 ± 85.69	545.79 ± 98.21	65.48	<.001	23.25	<.001	5.72	<.001
Week ₄	330.67 ± 50.40 ^{a,b}	358.65 ± 53.23 ^{a,c}	498.79 ± 60.27						
Week ₈	243.41 ± 49.19 ^{a,b}	261.39 ± 70.44 ^{a,c}	520.88 ± 63.87						
Week ₁₂	234.66 ± 56.06 ^{a,b}	279.98 ± 87.23 ^{a,c}	531.70 ± 63.92						
Week ₁₆	194.70 ± 83.76 ^{a,b}	227.52 ± 113.06 ^{a,c}	558.55 ± 90.29						
Area (cm ²)									
Week ₀	2.50 ± 1.15	2.28 ± 0.86	2.46 ± 0.91	5.50	.005	5.78	<.001	0.92	.502
Week ₄	2.37 ± 1.07	2.17 ± 1.04	2.28 ± 0.31						
Week ₈	1.43 ± 1.09 ^a	1.67 ± 1.01	2.33 ± 0.57						
Week ₁₂	1.40 ± 0.79 ^a	1.41 ± 0.82 ^a	2.15 ± 0.52						
Week ₁₆	1.14 ± 0.97 ^{a,b}	1.25 ± 0.94 ^a	2.17 ± 0.71						

(continued)

Table 3 (continued)

	TCC Group (n = 12)	BW Group (n = 13)	C Group (n = 11)	Group		Time		Group × Time	
				F	p	F	p	F	p
D-ap (mm)									
Week ₀	41.15 ± 6.12	40.07 ± 7.55	38.96 ± 9.50	20.16	<.001	12.76	<.001	3.24	.002
Week ₄	36.35 ± 8.37	37.06 ± 9.13	40.94 ± 11.05						
Week ₈	37.00 ± 8.20	28.76 ± 8.72 ^{a,c}	40.36 ± 7.57						
Week ₁₂	28.54 ± 7.42 ^{a,b}	24.52 ± 6.21 ^{a,c}	40.38 ± 8.84						
Week ₁₆	25.88 ± 8.95 ^{a,b}	21.54 ± 7.38 ^{a,c}	37.98 ± 7.02						
D-ml (mm)									
Week ₀	39.33 ± 8.85	41.75 ± 6.24	38.96 ± 9.50	21.02	<.001	13.01	<.001	3.24	.002
Week ₄	38.52 ± 11.89	41.99 ± 18.10	40.94 ± 11.05						
Week ₈	31.07 ± 7.92 ^a	39.24 ± 12.91	36.72 ± 5.54						
Week ₁₂	27.13 ± 8.15 ^a	35.51 ± 17.51	36.74 ± 9.74						
Week ₁₆	25.40 ± 9.83 ^a	30.52 ± 14.98 ^a	36.16 ± 8.72						

Note. Data are presented as mean ± SD. TCC = Tai Chi Chuan; BW = brisk walking; C = control; Time = the length of time with single-leg stance; Lng = the total length of the COP trajectories; Area = the area of 95% bivariate confidence ellipse, which encloses approximately 95% of the points on the COP path; D-ap = the maximal displacement of COP in the anterior-posterior direction; D-ml = the maximal displacement of COP in the medial-lateral direction; COP = center of pressure.

^aSignificant difference compared with the value at week₀ within each group. ^bSignificant difference between TCC and C groups. ^cSignificant difference between BW and C groups.

This finding is consistent with those of previous studies (Maciaszek & Osinski, 2012; Won, Kim, & Oh, 2015). A previous study (Zhou et al., 2015) found that a 24-week TCC exercise can significantly reduce total, AP, and ML COP sway paths in elderly women. In another study, postural stability with eyes closed was significantly improved after a 10-week BW intervention among the elderly (Gába et al., 2016), whereas in the current work, significant improvement was found after 12 weeks. The difference in age (63.2 vs. 55.9 years) might lead to differences in testing results. Another study (Li, Xu, & Hong, 2008) showed that a 16-week intervention failed to improve the testing scores for SLC. The difference in training frequency might be another factor leading to difference in testing results.

Several factors may improve balance ability after regular TCC and BW interventions. Muscle strength, proprioception, and the neuromuscular reaction ability of low extremities play important roles in controlling body balance (Jacobson, Chen, Cashel, & Guerrero, 1997). TCC intervention reportedly increases the muscle strength of knee flexors and extensors (Xu et al., 2008) and could decrease neuromuscular reaction time in the rectus femoris/semitendinosus/anterior tibialis/gastrocnemius muscles (Sun et al., 2016). Regular TCC and BW interventions could also decrease ankle kinesthesia threshold (Zhang et al., 2014).

Our data indicated that 12-week TCC and 16-week BW exercises could improve balance ability, respectively. To enhance body balance for preventing falls, the elderly must persist in doing exercises for several months. Moreover, TCC exercise may be more efficient than BW in improving balance ability with SLO (TCC after 8 weeks and BW after 12 weeks) and SLC (TCC after 12 weeks and BW after 16 weeks). The more efficient improvement of balance ability in the TCC group than in the BW group could be attributed to the characteristics of the movements involved. The TCC exercise emphasizes a large shift of the center of mass, long durations of SLS, large ankle and knee joint range of motion, and repeated concentric and eccentric contractions of lower muscles (Mao, Hong, & Li, 2006). In addition, a study indicated that the greatest pressure–time integral regions during TCC movements are the great toe and first metatarsal head. By contrast, the greatest pressure–time integral regions during walking are the second and third metatarsal heads (Mao, Li, & Hong, 2006). The cutaneous afferent information from the great toe plays an important role in postural control (Tanaka, Takeda, Izumi, Ino, & Ifukube, 1999). A study found a negative relation between plantar pressure and cutaneous sensation threshold of the great toe (Nurse & Nigg, 2001) from cutaneous mechanoreceptors located on the plantar of the feet, which could detect pressure and skin deformations and provide somatosensory information for the body (Yümin, Şimşek, Sertel, & Ankaralı, 2016). Thus, when the plantar pressure in the great toe is high, the cutaneous sensation is highly sensitive.

Our study showed that the periods to improve balance under SLO were shorter than that under SLC (8 vs. 12 weeks). Vision is one of the most important information input systems contributing to postural control. Clear and precise vision could provide the accurate and necessary spatial location information to the central nervous system. This information on upright posture serves as effective feedback in moderating balance control to prevent falls (Sturnieks, George, & Lord, 2008). Compared with that in the eyes-open condition, the COP sway area under the eyes-closed condition increased at approximately 30% while standing on a firm surface

(Ahead of Print)

(Lord, Clark, & Webster, 1991). Postural control ability under SLC could be more difficult than that under SLO. Thus, long periods maybe necessary to improve balance ability under the eyes-closed condition. The authors speculate that visual conditions could be the main reason for the different periods necessary to improve balance ability under such visual conditions.

Our data also showed that balance control improved better in the AP direction than that in the ML direction in the BW group. BW exercise enhanced balance control more than the TCC did in the AP direction. First, BW was more effective in decreasing maximal displacements with SLC in the AP direction (after 8 weeks) than in the ML direction (after 16 weeks). The differences can be associated with the information input system. The standing balance control sway without vision relied on the joint proprioceptive feedback and increased by 20–70% relative to that under visual feedback (Lord & Dayhew, 2001). The kinesthesia threshold of the plantar–dorsal flexion in the AP direction was significantly lower than that of inversion–eversion in the ML direction (Zhang et al., 2014). Moreover, a study found that the ankle joint proprioception in the ML direction failed to significantly improve during a 16-week BW exercise program (Zhang et al., 2014). Second, in our study, the D-ap in the BW group with SLC significantly decreased earlier (TCC vs. BW, 12 weeks vs. 8 weeks) than the TCC group. These results indicated that the balance improvements in the AP direction of the BW group were more efficient than that of the TCC group. The different movement characteristics between the TCC and BW exercises might lead to a discrepancy in intervention effects on postural control. Different postural control mechanisms were found in the AP direction compared with those in the ML direction. Balance stability was primarily controlled by the “ankle joint strategy” in the AP direction and the “hip joint strategy” in the ML direction (Corrigan, Cashman, & Brady, 1992). Walking movements included the ankle/knee joints flexed and repeatedly extended in the sagittal plane. However, ankle joint movement was multidirectional in the TCC exercise. This special uniaxial movement characteristic of BW may help enhance the musculoskeletal system function (muscle strength, neuromuscular reaction, and proprioception) in the AP direction.

The present study has two limitations. First, only female participants were recruited. Thus, further study should explore the effects of TCC and BW on the balance ability of elderly men. Second, only 36 participants completed the entire 16-week study, so the findings of this study should be interpreted with caution. Further studies with large sample sizes should be required to determine the effects of TCC and BW interventions on balance in elderly.

Conclusion

The 12-week TCC and 16-week BW interventions could improve the balance ability of elderly women. Compared with BW, TCC is more efficient in improving the body balance ability under SLS. Moreover, improving balance ability under SLS is quicker with vision than without vision. Thus, adequate periods of TCC and BW exercises are essential to improve the balance ability of elderly women, especially with their eyes closed.

(Ahead of Print)

Acknowledgment

This work was supported by the National Natural Science Fund of China (31700815, 11572202) and the Shandong Science & Technology Department (2017G006043).

References

- Chan, A.W., Sit, J.W., Chair, S.Y., Leung, D.Y., Lee, D.T., Wong, E.M., & Fung, L.C. (2016). Evaluation of the effectiveness of Tai Chi versus brisk walking in reducing cardiovascular risk factors: Protocol for a randomized controlled trial. *International Journal of Environmental Research and Public Health*, 13(7), 682. doi:10.3390/ijerph13070682
- Chang, S., Zhou, J., Hong, Y., Sun, W., Cong, Y., Qin, M., ... Li, W. (2016). Effects of 24-week Tai Chi exercise on the knee and ankle proprioception of older women. *Research in Sports Medicine*, 24(1), 84–93. doi:10.1080/15438627.2015.1126281
- Corrigan, J.P., Cashman, W.F., & Brady, M.P. (1992). Proprioception in the cruciate deficient knee. *The Journal of Bone & Joint Surgery*, 74, 247–250. PubMed doi:10.1302/0301-620X.74B2.1544962
- Davis, J.C., Robertson, M.C., Ashe, M.C., Liu-Ambrose, T., Khan, K.M., & Marra, C.A. (2010). International comparison of cost of falls in older adults living in the community: A systematic review. *Osteoporosis International*, 21(8), 1295–1306. doi:10.1007/s00198-009-1162-0
- Dixit, S., Maiya, A., Shastry, B.A., & Guddattu, V. (2016). Analysis of postural control during quiet standing in a population with diabetic peripheral neuropathy undergoing moderate intensity aerobic exercise training: A single blind, randomized controlled trial. *American Journal of Physical Medicine & Rehabilitation*, 95(7), 516–524. PubMed doi:10.1097/PHM.0000000000000426
- Enderlin, C., Rooker, J., Ball, S., Hippensteel, D., Alderman, J., Fisher, S.J., ... Jordan, K. (2015). Summary of factors contributing to falls in older adults and nursing implications. *Geriatric Nursing*, 36(5), 397–406. PubMed doi:10.1016/j.gerinurse.2015.08.006
- Franchignoni, F., Tesio, L., Martino, M.T., & Ricupero, C. (1998). Reliability of four simple, quantitative tests of balance and mobility in healthy elderly females. *Aging Clinical and Experimental Research*, 10(1), 26–31. doi:10.1007/BF03339630
- Gába, A., Cuberek, R., Svoboda, Z., Chmelík, F., Pelciová, J., Lehnert, M., & Frömel, K. (2016). The effect of brisk walking on postural stability, bone mineral density, body weight and composition in women over 50 years with a sedentary occupation: A randomized controlled trial. *BMC Women's Health*, 16(1), 1–10. doi:10.1186/s12905-015-0282-2
- Gerbino, P.G., Griffin, E.D., & Zurakowski, D. (2007). Comparison of standing balance between female collegiate dancers and soccer players. *Gait & Posture*, 26(4), 501–507. PubMed doi:10.1016/j.gaitpost.2006.11.205
- Gribble, P.A., Tucker, W.S., & White, P.A. (2006). Time-of-day influences on static and dynamic postural control. *Journal of Athletic Training*, 42(1), 35–41.
- Hijmans, J.M., Zijlstra, W., Geertzen, J.H., Hof, A.L., & Postema, K. (2009). Foot and ankle compression improves joint position sense but not bipedal stance in older people. *Gait & Posture*, 29(2), 322–325. doi:10.1016/j.gaitpost.2008.10.051
- Huang, Y., & Liu, X. (2015). Improvement of balance control ability and flexibility in the elderly Tai Chi Chuan (TCC) practitioners: A systematic review and meta-analysis. *Archives of Gerontology and Geriatrics*, 60(2), 233–238. PubMed doi:10.1016/j.archger.2014.10.016

(Ahead of Print)

- Jacobson, B.H., Chen, H., Cashel, C., & Guerrero, L. (1997). The effect of T'ai Chi Chuan training on balance, kinesthetic sense, and strength. *Perceptual and Motor Skills*, 84(1), 27–33. PubMed doi:10.2466/pms.1997.84.1.27
- Kathiresan, G., Jali, N., Afifah, N., Aznie, N., & Fideyana, N. (2010). The relationship between ankle joint flexibility and squatting knee flexion posture in young Malaysian men. *World Journal of Sport Sciences*, 3(3), 226–230.
- Kim, H., Kim, Y.L., & Lee, S.M. (2015). Effects of therapeutic Tai Chi on balance, gait, and quality of life in chronic stroke patients. *International Journal of Rehabilitation Research*, 38(2), 156–161. PubMed doi:10.1097/MRR.0000000000000103
- Konak, H.E., Kibar, S., & Ergin, E.S. (2016). The effect of single-task and dual-task balance exercise programs on balance performance in adults with osteoporosis: A randomized controlled preliminary trial. *Osteoporosis International*, 27(11), 3271–3278. doi:10.1007/s00198-016-3644-1
- Li, F., Harmer, P., Fitzgerald, K., Eckstrom, E., Stock, R., Galver, J., ... Batya, S.S. (2012). Tai Chi and postural stability in patients with Parkinson's disease. *New England Journal of Medicine*, 366(6), 511–519. PubMed doi:10.1056/NEJMoal107911
- Li, J., Xu, D., & Hong, Y. (2008). Effects of 16-week Tai Chi intervention on postural stability and proprioception of knee and ankle in older people. *Age and Ageing*, 37(5), 575–578. doi:10.1093/ageing/afn109
- Lord, S.R., Clark, R.D., & Webster, I.W. (1991). Postural stability and associated physiological factors in a population of aged persons. *Journal of Gerontology*, 46(3), M69–M76. doi:10.1093/geronj/46.3.M69
- Lord, S.R., & Dayhew, J. (2001). Visual risk factors for falls in older people. *Journal of the American Geriatrics Society*, 49(5), 508–515. PubMed doi:10.1046/j.1532-5415.2001.49107.x
- Maciaszek, J., & Osinski, W. (2012). Effect of Tai Chi on body balance: Randomized controlled trial in elderly men with dizziness. *The American Journal of Chinese Medicine*, 40(2), 245–253. PubMed doi:10.1142/S0192415X1250019X
- Mao, D., Hong, Y., & Li, J. (2006). Characteristics of foot movement in Tai Chi exercise. *Physical Therapy*, 86(2), 215–222. PubMed
- Mao, D., Li, J., & Hong, Y. (2006). Plantar pressure distribution during Tai Chi exercise. *Archives of Physical Medicine and Rehabilitation*, 87(6), 814–820. PubMed doi:10.1016/j.apmr.2006.02.035
- Murtagh, E.M., Boreham, C.A., & Murphy, M.H. (2002). Speed and exercise intensity of recreational walkers. *Preventive Medicine*, 35(4), 397–400. PubMed doi:10.1006/pmed.2002.1090
- Nilsson, G., Ageberg, E., Ekdahl, C., & Eneroth, M. (2006). Balance in single-limb stance after surgically treated ankle fractures: A 14-month follow-up. *BMC Musculoskeletal Disorders*, 7(1), 1–8. doi:10.1186/1471-2474-7-35
- Novak, A.C., Komisar, V., Maki, B.E., & Fernie, G.R. (2016). Age-related differences in dynamic balance control during stair descent and effect of varying step geometry. *Applied Ergonomics*, 52, 275–284. PubMed doi:10.1016/j.apergo.2015.07.027
- Nurse, M.A., & Nigg, B.M. (2001). The effect of changes in foot sensation on plantar pressure and muscle activity. *Clinical Biomechanics*, 16(9), 719–727. PubMed doi:10.1016/S0268-0033(01)00090-0
- Okubo, Y., Osuka, Y., Jung, S., Rafael, F., Tsujimoto, T., Aiba, T., ... Tanaka, K. (2015). Walking can be more effective than balance training in fall prevention among community-dwelling older adults. *Geriatrics & Gerontology International*, 16(1), 1–8.
- Paillard, T., Lafont, C., Costes-Salon, M.C., Rivière, D., & Dupui, P. (2004). Effects of brisk walking on static and dynamic balance, locomotion, body composition, and aerobic capacity in ageing healthy active men. *International Journal of Sports Medicine*, 25(7), 539–546. PubMed doi:10.1055/s-2004-820948

(Ahead of Print)

Received: 31 August 2020 | Revised: 22 October 2020 | Accepted: 31 October 2020
DOI: 10.1111/jpc.12884

ORIGINAL ARTICLE

WILEY

The effectiveness of Tai Chi Chuan on fear of movement, prevention of falls, physical activity, and cognitive status in older adults with mild cognitive impairment: A randomized controlled trial

Canan Birimoglu Okuyan PhD¹ | Ebru Deveci MSc²

¹Department of Public Health Nursing, Faculty of Health Sciences, Sakarya University of Applied Sciences, Sakarya, Turkey

²Department of Anesthesia, Vocational School of Health Services, University of Hasan Kalyoncu, Gaziantep, Turkey

Correspondence

Canan Birimoglu Okuyan, Division of Gait and Balance, University of Western Ontario, London, Ontario N6C 6A7, Canada.

Email: cananbirimoglu@gmail.com, cananbirimoglu@uwo.ca and cbrimog@uwo.ca

Abstract

Purpose: This study aimed to investigate the effectiveness of Tai Chi Chuan (TCC) on fear of movement, prevention of falls, physical activity, and cognitive status in older adults with mild cognitive impairment.

Design and Methods: This controlled trial was conducted with 20 participants in the intervention group, and they did exercises for 35–40 min/session, twice a week for 12 weeks.

Finding: The posttest scores of the scales indicated that the scores of cognitive adaptations, being observant, and changes in level subscales of the fall behavioral scale in the older adults who did Tai Chi exercise in the intervention group significantly increased ($p < .01$).

Practice Implications: The nursing home provider can safely manage TCC exercise interventions for older adults with MCI.

KEYWORDS

mild cognitive impairment, older adults, risk of falling, Tai chi chuan

1 | INTRODUCTION

Mild cognitive impairment (MCI) is being a common disorder experienced by elderly individuals and also a significant public health problem.¹ MCI is regarded as a transient stage between healthy aging and dementia and is characterized by memory problems and the decline of cognitive functions.² The impaired cognitive functions cause a significant decrease in the quality of life, activities of daily living, and independence of individuals with MCI.³ Also, elderly individuals with MCI have a higher risk of dementia and falls, as well as low balance.^{4,5}

There has been no effective treatment yet to alleviate the clinical outcomes of MCI.⁶ Therefore, there is a considerable need to develop preventive interventions and therapeutic approaches to prevent the progression of dementia.⁷ Recent studies on nonpharmacological interventions have suggested that lifestyle factors, such as physical activity may contribute to modifying disease progression.^{8–10} Also,

decreased levels of physical activity may be a risk factor for dementia in older adults. The previous study has highlighted the importance of physical activity programs that are effective in fear of movement, prevention of falls, and improving cognitive functions.¹¹ However, no consensus has yet been reached regarding exercises to be given to patients with MCI.¹² It has been stated that the most appropriate exercise training program that can be given to MCI individuals should include parameters, such as balance, flexibility, lower and upper extremity muscle strengthening exercises, and goal-oriented activities.¹³

Tai Chi Chuan (TCC), one of the aerobic exercises maintaining muscle strength and balance, is the most recommended traditional Chinese exercise for older adults.¹⁴ TCC diminishes the risk of falls, ameliorates balance, and enhances endurance and flexibility of muscle.^{15,16} The current evidence has revealed that Tai Chi can be considered as an ideal exercise for older adults to improve cognitive functions and quality of life.¹⁴

As long as the value of improving cognitive health and preventing falls in the elderly has increased worldwide, MCI has attracted significant attention from researchers. However, there has been no study examining the effectiveness of exercise programs, such as TCC on fear of movement, prevention of falls and cognition in people with MCI in the Turkish population. Furthermore, A research study has highlighted that there are physical inactivity and low levels of regular exercise in older adults in the Turkish population.²² Therefore, it is incredibly crucial to understand better the role of Tai Chi exercise in cognitive and physical health in the Turkish population. This study aimed to investigate the effectiveness of TCC on fear of movement, prevention of falls, physical activity, and cognitive status in older adults with MCI.

Hypotheses of the study

- (1) Is TCC exercise effects on the prevention of falls in older adults with MCI in the Turkish population?
- (2) Is TCC exercise effective on physical activity and fear of movement in older adults with MCI?
- (3) Is TCC exercise effective on cognitive adaptations in older adults with MCI?

2 | METHODS

2.1 | Study design

The present study was a randomized control trial conducted with pretest, posttest, and control-group design.

2.2 | Sample and setting

This study was conducted in a nursing home located in Turkey. Seventy-nine older adults live in this nursing home. The mini-mental state examination (MMSE),²³ Montreal cognitive assessment (MoCA)²⁴ tests, and clinical dementia rating scale (Stage 0.5–1)²⁵ were used to determine whether or not the older adults had MCI. The inclusion criteria of the study were determined as follows: (1) having a score of MMSE and MoCA of <25 points, (2) being Turkish and aged 65 years and over, (3) being able to communicate, read, and write independently in their local language, (4) being able to perform Tai Chi physical activity, (5) having no diagnosis of dementia and other psychiatric disorders. The exclusion criteria of the study were determined as follows: (1) having a body-mass index of >30, (2) having infectious diseases or immunological disorders, (3) being users of medications that would affect cognitive functions, (4) not signing the informed consent form.

The sample size of the present study was determined to be 34 (17 per group) with a power of 90% and Type I error of 5% using the power analysis (the program G*Power 3.1 was used). Considering the withdrawal possibility of 15% of participants, we enrolled 47 [control group = 24; intervention group = 23] older adults who met the

inclusion and exclusion criteria in this study. All participants were randomly assigned to the intervention (TCC) and control (untrained) groups. Three participants from the intervention group were excluded from the study since they could not continue the 12-week exercise program regularly. Also, two participants from the control group were excluded from the study since one of them refused to participate in the posttest, and the other was on leave for a long time. The data obtained from 42 participants who completed the study were analyzed. The individuals were stratified based on the age (between 65 and 74 years or being aged >75 years), gender (female or male) and the history of falling (yes or no) by using the blocked randomization method²⁶ and assigned to the intervention (TCC exercise) group ($n = 23$) and the control (untrained) group ($n = 24$) by drawing the lot. There was no significant difference between these groups ($p > .05$). The flow diagram of the Tai Chi Chuan exercise trial was shown in Figure 1.

2.3 | Intervention

The data were collected between October 2018 and December 2019. The personal information form, the Tinetti assessment tool (TAT), the Tampa scale of kinesiophobia (TSK), the physical activity scale for the elderly (PASE), and the falls behavioral (FaB) scale were applied as pretest by conducting the face-to-face interview with all the participants. After these forms were completed, patients in the intervention group started to do TCC exercises with the help of the researcher. As previously stated,¹² they performed the TCC exercise program every Wednesday and Thursday per week, like two sessions and 35 and 40-min sessions (Figure 2).

The PASE, TAT, FaB, and TSK had applied again as post-tests to the intervention group ($n = 20$) at the end of 12 weeks. The individuals in the control group ($n = 22$) were not subjected to any physical practice, and only the data collection forms (pretest and posttests) were applied and evaluated together with the intervention group.

2.4 | Outcome measures

The outcomes were evaluated before the TCC training and at the end of 12 weeks. The researcher was blind to all outcome evaluations to reduce bias. The data were collected using a personal information form, TAT, PASE, TSK, and FaB. Personal Information Form was prepared by the researcher upon the literature review.^{12,13} This form includes the life habits and health conditions of the participants.

TAT, which was developed by Mary Tinetti to evaluate the risk of falling in people with MCI, was used in the present study.¹⁷ The scale consists of a total of 28 items, including 12 items for gait and 16 items for balance. TAT was adapted to the Turkish population by Agrcan,²⁷ and it was found that the value of Cronbach's alpha coefficient was 0.97.²⁰ In the present study, Cronbach's alpha coefficient was found to be 0.96.




FIGURE 1 Flow diagram of the Tai Chi exercise trial. TCC, Tai Chi Chuan [Color figure can be viewed at wileyonlinelibrary.com]

PASE, which was developed in 1993 to assess the status of physical activity in people with MCI, was used in the present study. The PASE total score varies between 0 and 400, and higher scores signify high levels of physical activity.⁷¹ Ayva et al.²² adapted this scale to the Turkish population and determined that Cronbach's alpha coefficient of the scale was 0.71. In the present study, Cronbach's alpha coefficient was found to be 0.70.

TSK with 17 items was used to evaluate the fear of movement in the present study. The total score of TSK ranges from 17 to 68.⁷² Yilmaz et al. adapted TSK to the Turkish population and found that the test-retest reliability was 0.806.⁷³ In the present study, Cronbach's alpha coefficient was found to be 0.78.

The FoB scale, which was developed by Clemson et al., was used in the study to assess behaviors related to falls in people with MCI. The FoB scale includes 30 items and ten subscales. While higher



FIGURE 2 Example of Yang style of TCC. TCC, Tai Chi Chuan [Color figure can be viewed at wileyonlinelibrary.com]

scores indicate the safe or protective behaviors of the individuals related to falling, lower scores signify the risky behaviors.²⁵ Uymaz and Nalçivan adapted the F&B to the Turkish population and found that the Cronbach's alpha coefficient was 0.90.²⁶ In the present study, Cronbach's alpha coefficient was determined to be 0.89.

2.5 | Statistical analysis

The data of the study were analyzed through The statistical package for social sciences 22.0 version, and Graphs in the study were prepared via GraphPad Prism software. All data were represented as the mean $\pm$ SD. The Kolmogorov-Smirnov test and the Shapiro-Wilk test were used as the normality distribution test. The data regarding demographic and clinical characteristics in the intervention and control groups were analyzed by using χ^2 and t test. Differences in falls, balance, physical activity, and kinesiophobia were compared between intervention and control groups at both beginning and end of the study by using The Wilcoxon signed-rank test or paired t test where appropriate. The value of $p < .05$ was considered statistically significant.

2.6 | Ethical considerations

We obtained ethical approval (no: 2018/120) and institutional permissions before starting the study and conducted this study in accordance with the Declaration of Helsinki. We informed the participants about the current study and stated that the personal

identities of the participants will not be published and the participants had the freedom to withdraw from the study at any time without any reason being given. We obtained written informed consent from the elderly people before being included in the study.

3 | RESULTS

The mean age of geriatric individuals participating in the research was 74.21 ± 6.93 years. The 93.74% of the individuals in both groups were male in the experimental and control group (13 [56.5%] and 14 [58.3%], respectively), and the majority of older adults in the experimental and control groups had an education level of elementary school (56.5% and 45.8%, respectively). In both groups, the majority had social insurance (87.0% and 87.5%, respectively). Moreover, most participants in the experimental and control groups had balance issues (69.6% and 66.7%, respectively). Twenty-two males (mean age of 74.14 ± 7.8) and Twenty female (mean age of 74.30 ± 5.7) participants in the control and intervention groups completed the study. Seven participants from the intervention group could not complete the 12-week TCC training and thus were not included in the final analysis.

Tables 1 and 2 show the results of the present study, which was carried out to determine the effects of TCC as a mind-body movement therapy on the fear of movement, gait, risk of falling and cognitive improvement in older adults with MCI. The PASE, TBT & TGT, F&B, and TKS were applied as a pretest to all participants. It was determined in the present study that there was no significant difference between intervention and the control groups ($p > .05$).

TABLE 1 The mean pretest and posttest scores of the scales in the experimental group

Scales	Pretest		Posttest		The value and the significance of the test	
	<i>x</i>	<i>SD</i>	<i>x</i>	<i>SD</i>	<i>t</i>	<i>p</i>
Tinetti balance assessment	9.73	1.95	12.8	1.81	-10.788	.000
Tinetti gait assessment	10.6	0.88	11.7	0.64	-5.234	.000
TSK	42.94	3.31	48.21	3.5	-7.175	.000
PASE score (0-400)	41.77	12.10	62.27	13.82	-6.413	.000
Sub-dimensions scores of the FaB scale						
Cognitive adaptations	2.50	0.47	3.17	0.57	-3.042	.006
Protective mobility	2.48	0.50	2.86	0.62	-4.394	.000
Avoidance	2.38	0.60	2.39	0.61	-0.123	.903
Awareness	2.70	0.69	2.81	0.76	-1.796	.086
Hastiness	2.68	1.01	2.84	0.80	-1.567	.131
Practical strategies	2.00	1.08	2.08	1.04	-1.447	.162
Displacing activities	2.08	0.99	2.21	0.90	-1.817	.083
Being observant	2.04	0.76	2.43	0.72	-3.761	.001
Changes in level	2.82	0.38	3.39	0.49	-4.041	.001
Getting to the phone	2.73	0.81	2.78	0.67	-0.371	.714
Total of FaB scale	2.48	0.28	2.68	0.19	-5.010	.000

Abbreviations: FaB, falls behavioral; PASE, physical activity scale for the elderly; TSK, Tampa scale of kinesiophobia.

In the present study, a significant difference was observed in the pretest and posttest scores of the TAT, PASE, TSK, and FaB scale in the intervention group ($p < .05$). Also, the posttest scores of cognitive adaptations, protective mobility,

being observant, changes in level subscales of the FaB scale significantly increased in the intervention group ($p < .05$). On the other hand, the results of this study revealed that there was no significant difference between the pretest and posttest scores of

TABLE 2 The mean pretest and posttest scores of the scales in the control group

Scales	Pretest		Posttest		The value and the significance of the test	
	<i>x</i>	<i>SD</i>	<i>x</i>	<i>SD</i>	<i>t</i>	<i>p</i>
Tinetti balance assessment	9.95	1.73	9.45	1.66	1.906	.069
Tinetti gait assessment	8.50	2.26	8.54	2.24	-0.272	.788
TSK	43.16	3.5	43.37	3.5	-0.666	.512
PASE score (0-400)	42.86	13.01	42.70	13.27	0.335	.742
Sub-dimensions scores of the FaB scale						
Cognitive adaptations	2.76	0.43	2.75	0.53	0.115	.910
Protective mobility	2.47	0.49	2.43	0.56	1.000	.328
Avoidance	2.40	0.59	2.38	0.66	0.485	.632
Awareness	2.67	0.69	2.66	0.69	1.000	.328
Hastiness	2.65	1.00	2.61	0.57	0.296	.770
Practical strategies	1.95	0.75	1.91	0.97	0.272	.788
Displacing activities	2.08	0.97	1.95	0.90	1.141	.266
Being observant	2.20	0.78	2.16	0.76	0.225	.824
Changes in level	2.87	0.34	2.83	0.48	0.569	.575
Getting to the phone	2.79	0.78	2.75	0.73	1.000	.328
Total of FaB scale	2.48	0.28	2.44	0.23	1.307	.204

Abbreviations: FaB, falls behavioral; PASE, physical activity scale for the elderly; TSK, Tampa scale of kinesiophobia.




FIGURE 3 Comparison of posttest values of the scales in control and TCC groups. (A) Tinetti balance assessment, (B) Tinetti gait assessment, (C) Tampa scale for Kinesiophobia, (D) physical activity scale for elderly. * $p < .01$ compared with the control. All data were presented as mean $\pm$ SD. TCC, Tai Chi Chuan

the TAT, TSK, PASE, and subscales of the FaB scale in the control group ($p > .05$).

When TAT, TSK, and PASE posttest scores of the intervention and control groups were examined, all scores of these scales were significantly higher in the intervention group compared to the control group ($p < .01$) (Figure 3). Moreover, the posttest scores of cognitive adaptations, protective mobility, and changes in level subscales of the FaB scale were significantly higher in the intervention group than the control group (Figure 4) ($p < .01$).

It was found that TAT, TSK, and PASE scores of the intervention group significantly increased after the TCC exercise ($p < .01$). Likewise, the posttest scores of the scales revealed that scores of cognitive adaptations, protective mobility, being observant, and changes in level subscales of the FaB scale significantly increased in the older adults in the intervention group whom did TCC exercise ($p < .01$).

4 | DISCUSSION

The effects of 12-week TCC exercise on kinesiophobia, prevention of falls and cognitive functions in older adults with MCI were examined in this study. It was found that TCC exercise affected falls, physical activity levels, kinesiophobia, and falls-related safe or protective behaviors in older adults. Furthermore, it was

revealed that the TCC exercise positively affected these parameters, and the results of the study supported all three hypotheses. According to a previous study emphasized that aerobic exercise in individuals with MCI reduced cognitive impairment and prevention of falls in elderly individuals was also useful in terms of reducing the transition from MCI to dementia, and TCC was recommended as a new option in reducing the transition from MCI to dementia.²² However, the effect of TCC on individuals with MCI is not precisely elucidated.

The most common problems experienced by elderly individuals include fear of movement and balance disorders associated with falls, and these factors affect their daily activities and social relationships.^{25,29} It was observed that TAT scores of elderly adults who did TCC exercise in the intervention group increased; whereas, their fear of movement decreased. Moreover, an increase was observed in physical activity levels (PASE); on the other hand, a decrease was determined in safety or protective behavior related to falls (Table 1). These results suggested that older adults had a reduced risk of falls, increased safety and protective behaviors related to falls, and increased physical movement levels as a result of TCC training. Physical activities might have a positive effect on cognitive status in individuals with MCI. The previous study demonstrated that decreased physical performance in older adults with MCI might increase the risk of




FIGURE 4 The sub-dimensions scores of the FaB scale, such as (A) cognitive adaptations, (B) mobility, and (C) changes in level in the posttest values of the scales in control and TCC groups. * $p < .01$ compared with the control. All data were presented as mean $\pm$ SD. TCC, Tai Chi Chuan

Alzheimer's disease.²⁰ Therefore, increasing physical activity may prevent the development of cognitive impairment in older adults.^{21,22} Recent studies have also revealed that TCC is one of the best exercises to achieve this goal since balance can be developed with exercise that helps to improve joint movement and posture.^{22,23} It has been reported that older adults had improved balance and walking status after TCC exercise.^{22,23} Especially, Yang-style TCC is useful in terms of increasing musculoskeletal flexibility in healthy older women.²⁴ Balance exercises, such as TCC improve static and dynamic balance as well as performance in healthy older adults.¹⁵ In another study, it was observed that while resistance exercise did not improve posture control, balance exercises improved posture in older adults.²⁵ In their study, Sangalaj et al. divided 40 individuals with multiple sclerosis into four groups and applied aerobic and resistive exercises in different combinations to those groups. The groups continued their programs four days a week for eight weeks, and at the end of the study, more effective results were obtained in balance, strength, agility, fatigue, speed, and walking distance values in the aerobic exercise group.²⁷ Fear of movement can lead to the avoidance of physical movements and activities in older adults since they think that movement can cause an injury once again.²⁸ In the present study, the fear of movement was found to be less in the intervention group compared to the control group, suggesting that

regular exercise habits reduce the fear of movement in elderly individuals.

The present study has some limitations. It was a randomized controlled trial with a relatively small population, and the MCI diagnosis of participants in the intervention and control groups was determined using a mini-mental test (the cognitive status assessment with Iscore of MMSE and MoCA < 25 points) and clinical dementia rating scale (Stage 0.5–1).

5 | CONCLUSION

The results of the present randomized controlled trial suggested that TCC was effective in decreasing fear of movement and risk of falls and improving cognitive functions. TCC, which is not a challenging and high-speed exercise, is an appropriate exercise for elderly individuals. Furthermore, TCC exercise may be useful in reducing the transition to dementia and improving cognitive functions in patients with MCI. However, further comprehensive studies are required better to understand the effect of TCC in individuals with MCI. It is essential to increase the knowledge and awareness of healthcare professionals about the importance of TCC aerobic exercises in reducing the fear of falling in people with MCI. TCC aerobic exercises into the care practices of older adults.

Also, biochemical studies are needed to support the effectiveness of TCC in people with MCI.

5.1 | Implications for psychiatric nursing practice

The implications of this study suggest that it is exceptionally crucial to nurses' intentions to exercise with older adults in their professional practice in the Turkish population. Also, as far as we know, there is no study investigating the effect of TCC on falls, physical activity, and cognitive status and gait parameters in individuals with MCI in Turkey.

Health care providers should be knowledgeable about the TCC to provide sufficient information to individuals and should teach the elderly different types of physical activity they can perform. Also, care providers and caregivers can safely manage TCC exercise interventions for older adults with MCI. Ongoing support and exercise will promote older adults' commitment to their TCC exercise. Even though the majority of older adults should exercise in nursing homes, older adults in Turkey do not prefer to exercise. The addition of this TCC will interventions to patients with MCI enable nurses to adjust their nursing interventions to the needs of these patient populations. That is why it is crucial to increase the knowledge and awareness of healthcare professionals about the importance of TCC aerobic exercises in reducing the fear of falling, prevention of falls and improving cognitive functions in people with MCI. This article provides new information regarding the reduced risk of falls, sustain mental health, reducing the transition to dementia and improving cognitive functions in patients with MCI, increased safety and protective behaviors related to falls and increased physical movement levels as a result of TCC training. Also, the paper provides recommendations for nurses and other clinicians to assist in the implementation of TCC interventions to patients with MCI.

CONFLICT OF INTERESTS

The authors declare that there are no conflict of interest.

AUTHOR CONTRIBUTIONS

As the authors of this paper, they have declared that they all have made a substantial contribution to the information or material submitted for the publication and they have approved the final version of this manuscript. C.B.O. conceived and designed the study. C.B.O. and E.D. collected data. C.B.O. and E.D. wrote/drafted/edited the manuscript and interpreted the results. C.B.O. and E.D. conducted analyses, prepared graphs/figures and revised the manuscript. All authors approved the content of this manuscript.

ETHICS STATEMENT

Ethical approval from the Clinical Trials Ethics Committee of Medical Faculty of Hatay Mustafa Kemal University (2018/120) and official permission from The Ministry of Family and Social Policies of Hatay were obtained.

ORCID

Canan Birimoglu Okuyan <https://orcid.org/0000-0002-7339-6072>

Ebru Döveci <https://orcid.org/0000-0003-0166-4741>

REFERENCES

- Eshkoor SA, Hamid TA, Mun CY, Ng OK. Mild cognitive impairment and its management in older people. *Clin Interv Aging*. 2015;10:687-693. <https://doi.org/10.2147/CIA.S73922>
- Tangalos EG, Peterson RC. Mild cognitive impairment in geriatrics. *Clin Geriatr Med*. 2018;34(4):563-589. <https://doi.org/10.1016/j.cger.2018.06.005>
- Kang Y, Lee E. Quality of life and its factors in Korean elderly with mild cognitive impairment. *Clin Nurs Res*. 2018;27(7):871-889. <https://doi.org/10.1177/1054773817714561>
- Montero Odasso M, Speechley M. Falls in cognitively impaired older adults: implications for risk assessment and prevention. *J Am Geriatr Soc*. 2018;66(2):367-375. <https://doi.org/10.1111/jgs.15219>
- Zhang W, Liaw LF, Schwank M, Mills N, Gwynn JD, Clement L. Review of gait, cognition, and fall risks with implications for fall prevention in older adults with dementia. *Dement Geriatr Cogn Disord*. 2019;48(1-2):17-29. <https://doi.org/10.1159/000504940>
- Nazam H, Stikel A, Corona M, Zeller M, Melrose RJ, Wilkins SS. Potential Benefits of Physical Activity in MCI and Dementia. *Behav Neurol*. 2020;2020:7807856-10. <https://doi.org/10.1155/2020/7807856>
- Rakshi G, Sato ST, Akoupolos GS, Zarous AS. Strategies for dementia prevention: latest evidence and implications. *Ther Adv Chronic Dis*. 2017;8(8):121-136. <https://doi.org/10.1177/2040622317712442>
- Northey JM, Cherbuin N, Pumpa KL, Smee DJ, Rattray B. Exercise interventions for cognitive function in adults older than 50: a systematic review with meta-analysis. *Br J Sports Med*. 2018;52(3):154-160. <https://doi.org/10.1136/bjsports-2016-096587>
- Munichowski P, Torosa M, Kubicki A, Mouray F. Impact of ambulatory physiotherapy on motor abilities of elderly subjects with Alzheimer's disease. *Geriatr Gerontol Int*. 2014;14(1):167-175. <https://doi.org/10.1111/ggi.12075>
- Bisbe M, Fuente Vidal A, López E, et al. Comparative cognitive effects of choreographed exercise and multimodal physical therapy in older adults with amnesia, mild cognitive impairment: randomised clinical trial. *J Alzheimers Dis*. 2020;73(2):769-783. <https://doi.org/10.3233/JAD-190552>
- Jia X, Jiang C, Tao J, Li Y, Zhou Y, Chen LD. Effects of core strength training combined with Tai Chi Chuan for the musculoskeletal system and cardiopulmonary function in older adults: A study protocol for a randomized controlled trial. *Medicine*. 2018;97(35):e12024. <https://doi.org/10.1097/MD.00000000000012024>
- Hosseini L, Karguzar E, Shari F, Negrindeh R, Memari AH, Nasab E. Tai Chi Chuan can improve balance and reduce fear of falling in community dwelling older adults: a randomized control trial. *J Exerc Rehabil*. 2018;14(6):1024-1031. <https://doi.org/10.12965/jer.1806024>
- Birimoglu Okuyan C, Bilgili N. Effect of Tai Chi Chuan on fear of falling, balance, and physical self-perception in elderly: a randomised controlled trial. *Turkish Journal of Geriatrics*. 2017;20(3):232-241.
- Siu MY, Lee DTF. Effects of Tai Chi on cognition and instrumental activities of daily living in community dwelling older people with mild cognitive impairment. *BMC Geriatr*. 2018;18(1):37. <https://doi.org/10.1186/s12877-018-0720-6>
- Folstein MF, Folstein SE, McHugh PR. "Mini-mental state": A practical method for grading the clinician. *J Psychiatr Res*. 1975;12(3):129-138. [https://doi.org/10.1016/0022-3956\(75\)90066-6](https://doi.org/10.1016/0022-3956(75)90066-6)
- Nasreddine ZS, Phillips NA, Bedirian V, et al. The montreal cognitive assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc*. 2005;53(9):695-699. <https://doi.org/10.1111/j.1532-5415.2005.03221.x>

THE BENEFITS OF TAI-CHI EXERCISE ON BALANCE CONTROL IN ELDERLY DURING STAIR-TO-FLOOR TRANSITION

Chenfu Huang and Tzu-hsiang Yang

Department of Physical Education, National Taiwan Normal University, Taipei, Taiwan

The purpose of this study is to investigate the benefits of Tai-Chi Exercise on balance control in elderly during the stair-to-floor transition. Twelve Tai-Chi elder participants and 12 normal healthy elderly were recruited. A Kistler force plate (1000Hz) obtained the COP (center of pressure) data and kinematics calculated from ten Vicon high-speed cameras (250Hz). Results revealed that Tai-Chi group descended at a faster speed and had larger step length. Tai-Chi group also had larger A-P (anteroposterior) COM-COP separation, A-P COP path and faster COP average velocity. The results indicated that long-term Tai-Chi exercise can improve the balance in elderly during the stair-to-floor transition.

KEYWORDS: center of pressure, step length.

INTRODUCTION: Falls were the leading cause of injuries in elderly (CDC, 2014), and fatal injuries occurred during stair walking. Falls during the descent of stair negotiation occur at least three times more frequently than those associated with stair ascent (Svanstrom, 1974). The stair-to-floor transition is the unique anticipation of the upcoming surface between stair and level walking (Sheehan & Gottschall, 2011), it would be higher fall risks than stair and level walking.

The change of center of pressure (COP) reflects the response of the central nervous system to the movement of the body's center of mass (COM), and the displacement of COP is used as a key factor of posture control and balance (Mao, Li, & Hong, 2006a). To modulate COP within the appropriate range of COM-COP separation is the key to maintaining dynamic stability and avoiding falls. To avoid falls, elderly adopted safer strategies while stair descent, including smaller A-P (anteroposterior) and M-L (mediolateral) COP displacement and slower COP movements (Kim, 2009a). During level walking and crossing obstacles, elderly reduced the A-P COM-COP separation to maintain balance (Hahn & Chou, 2004; Scarborough, Krebs, & Harris, 1999). However, the A-P distance is limited by the width of the ladder during stair descent, A-P COM-COP separation was no difference between the elderly and the young adults (Mian et al., 2007).

Tai-Chi is a popular traditional Chinese exercise for elderly, and the slow-motion characteristics of movement are quite suitable for older people. Tai-Chi gait emphasizes lowering the center of gravity and larger steps, showed more challenge to the balance ability and the lower limb muscles than the normal gait (Wu, 2008), and large A-P COP movement (Mao, Li, et al., 2006a). Tai-Chi exercise also improved the dynamic balance, increased the balance of single leg stand (Mao, Li, & Hong, 2006b) and reduced the fear of falling (Logghe et al., 2010). Even while the single-legged jumping and single-legged landing, the Tai-Chi elderly had better performance (Gyllenstein, Hui-Chan, & Tsang, 2010). It indicated that Tai-Chi exercise improved the dynamic balance of elderly, and it might also affect the performance of the daily life. Therefore, the purpose of this study is to investigate the benefits of Tai-Chi exercise on balance control in elderly during the stair-to-floor transition.

METHODS:

Twelve elderly that were regular Tai-Chi practitioners (eight males and four females) (Tai-Chi group), and twelve healthy elderly (eight males and four females) (normal group) participated in this study. Exclusion criteria were any known neurological or orthopedic disease and any current difficulties impeding their typical locomotion. Ethical approval was obtained from the Joint Institutional Review Board at Taipei Medical University. All participants provided their written informed consent before participation in our investigation. Tai-Chi group practiced

Yang style Tai-Chi which included 108 different body postures for an average of 10 years with a range between 2 to 15 years, four to five days per week, for a minimum of one hour per session. Control group of elderly performed leisure exercises (e.g. jogging, swimming) approximately over the same period per session but did not regularly perform Tai-Chi.

A ten camera motion system sampling at 250Hz and one Kistler force plate sampling at 1000Hz were synchronized to collect three-dimensional body motion and COP data. The 46 cm staircase consisted of a series of three steps, each step with a rise 18 cm and a run of 28 cm. Participants walked down the stairs starting with their dominant (right) leg with barefoot at a self-selected speed in a step-over-step manner and ended after the participant walked forward on level ground for approximately 5 m. The speed and step lengths were not restricted to simulate their typical performance. All participants completed at least five successful trials where the entire lead foot had contact with the force plate.

The stair-to-floor transition analysis began with toe-off of the right leg from the middle step (36cm height) to toe-off of the right leg from ground level. The COM velocity was calculated by averaging the 3-dimensional velocity of COM. Step length was defined as the anteroposterior distance between right toe and left toe during right leg ground contact and right leg toe-off the ground (shown in figure 1). An independent samples t-test was used to examine differences between groups. The alpha level (α) was set to 0.05 and all statistical analyses were conducted within SPSS 21.0.

RESULTS: The information of participants are shown in table 1. There were no significant differences in body weight and height, except in age. Tai-Chi group were older than the normal group.

Table 1: Information of participants.

	Tai-Chi group	Normal group	<i>p</i>
Age (years)*	72.7 $\pm$ 5.1	67.5 $\pm$ 2.7	<.01
Height (m)	1.63 $\pm$ 0.07	1.62 $\pm$ 0.06	
Weight (kg)	57.9 $\pm$ 0.07	63.5 $\pm$ 5.7	

The data of COP, temporal and kinematic variables are shown in table 2. Tai-Chi group descended at a faster speed and had larger step length. Tai-Chi group also had larger A-P COM-COP separation, M-L COP displacement and faster COP average velocity.

Table 2: COP, temporal and kinematic variables.

	Tai-Chi group	Normal group	<i>p</i>
A-P COM-COP separation			
1 st peak (cm)*	24.8 $\pm$ 3.5	21.4 $\pm$ 3.4	.02
2 nd peak (cm)*	35.1 $\pm$ 2.6	30.2 $\pm$ 4.0	<.01
M-L COM-COP separation (cm)	12.3 $\pm$ 2.6	11.3 $\pm$ 2.6	.32
A-P COP path (cm)	11.6 $\pm$ 2.7	10.2 $\pm$ 2.0	.16
M-L COP path (cm)*	10.7 $\pm$ 3.5	7.0 $\pm$ 1.5	<.01
COP average velocity (cm/s)*	100.2 $\pm$ 66.1	56.7 $\pm$ 14.3	.04
Stance time (s)	0.66 $\pm$ 0.08	0.68 $\pm$ 0.10	
1 st step length (m)*	0.51 $\pm$ 0.07	0.41 $\pm$ 0.07	<.01
2 nd step length (m)*	0.58 $\pm$ 0.04	0.53 $\pm$ 0.05	.02
COM velocity (m/s)*	0.93 $\pm$ 0.10	0.77 $\pm$ 0.12	<.01




Figure 1. 1st step length and 2nd step length during the stair-to-floor transition.

DISCUSSION: Modulating the translation between COM and COP is an important factor to maintain dynamic stability. The backward displacement of the COP can generate the momentum for forwarding motion (Polcyn, Lipsitz, Kerrigan, & Collins, 1998). Because of weakness in lower extremities muscles, elderly would reduce forward motion of COM and backward displacement of COP. After Tai-Chi exercise intervention, COP displacement increased while level walking, crossing obstacles and descending stairs (Kim, 2009b; Kim, Han, & Cho, 2009). The results showed that Tai-Chi group had larger A-P COP displacement, which is consistent with the previous studies. To simulate the performance in daily life, the study didn't restrict the descending speed and step length. Tai-Chi group had faster speed during stair descent, and a larger step length could modulate the forward rotational momentum (van Dieën, Spanjaard, Konemann, Bron, & Pijnappels, 2007). During gait, greater A-P COM-COP separation provided data that support more mechanic loading and improve ability to tolerate unsteadiness after Tai-Chi training (Gatts & Woollacott, 2007), therefore, older adults used a compensatory strategy by reducing the A-P COM-COP separation to maintain balance (Hahn & Chou, 2004; Scarborough et al., 1999). In this study, the larger 1st peak of A-P COM-COP separation is due to greater step length in Tai-Chi group. It indicated that Tai-Chi group had better ability to maintain dynamic stability during contact ground. At the end of the stair-to-floor transition, Tai-Chi group showed larger 2nd step length and the 2nd peak of A-P COM-COP separation, which COM was front from COP. Due to faster COM velocity and similar stance time, larger A-P COM-COP separation represented Tai-Chi exercise could increase COM path. (Gatts & Woollacott, 2007). In the frontal plane, Tai Chi exercise could increase the M-L COP path in elderly while contact ground (Kim et al., 2009), and increased M-L COP path indicated an improvement of coordination between hip adduction and abduction muscles after Tai-Chi training (Winter, Patla, Ishac, & Gage, 2003). During stair descent, the hip abductor muscles of the contralateral leg shifted COP to lateral side, and then the hip joint and the ankle joint muscles propelled COM forward until contralateral leg contact ground, therefore, the muscular coordination of frontal hip joint muscles is important for the elderly to maintain lateral stability during the stair-to-floor transition. The study showed that Tai-Chi group had greater M-L COP path, which indicated that Tai-Chi group had better lateral stability during contact ground. The COP average velocity provided important information on how to modulate the steps when facing challenges, such as stairs (Reid, Novak, Brouwer, & Costigan, 2011). Slower COP average velocity in elderly than that in young adults, because the slower COP average velocity is easier to maintain postural stability (Kim, 2009a; Reid et al., 2011). Previous study indicated COP average velocities were about 102~145cm/s and 58~85cm/s in young adults and elderly, respectively (Kim, 2009b). The results of the study showed that the Tai-Chi group had a faster COP average velocity, and it resembled the performance of young adults, which might reflect that Tai-Chi exercise improved the ability to maintain dynamic balance. The age of all participants in the study was above 65 years or older, the muscular power and work decreased by an average of 6% per decade (Daubney & Culham, 1999). In this study

Tai-Chi group were older than normal group, the better dynamic performance and balance control of Tai-Chi group indicated the benefits of long-term practicing Tai-Chi exercise on dynamic performance and balance control during stair-to-floor transition than the normal elderly.

CONCLUSION: Long-term Tai Chi exercise participant could improve the balance in elderly during the stair-to-floor transition. It is recommended that participation in Tai-Chi exercise will improve the balance and performance in daily activities.

REFERENCES

- CDC (Centers for Disease Control and Prevention). (2014). *Falls are leading cause of injury and death in older Americans*. Retrieved from <https://www.cdc.gov/media/releases/2016/p0922-older-adult-falls.html>
- Daubney, M. E., & Culham, E. G. (1999). Lower-extremity muscle force and balance performance in adults aged 65 years and older. *Physical therapy*, 79(12), 1177-1185.
- Gatts, S. K., & Woodlucott, M. H. (2007). How Tai Chi improves balance: biomechanics of recovery to a walking slip in impaired seniors. *Gait and Posture*, 25(2), 205-214.
- Gyllenstein, A. L., Hui-Chan, C. W., & Tsang, W. W. (2010). Stability limits, single-leg jump, and body awareness in older Tai Chi practitioners. *Arch Phys Med Rehabil*, 91(2), 215-220.
- Hahn, M. E., & Chou, L. S. (2004). Age-related reduction in sagittal plane center of mass motion during obstacle crossing. *Journal of biomechanics*, 37(6), 837-844.
- Kim, H. D. (2009a). A comparison of the center of pressure during stair descent in young and healthy elderly adults. *Journal of Physical Therapy Science*, 21(2), 129-134.
- Kim, H. D. (2009b). Age-related changes in the center of pressure trajectory during obstacle crossing. *Journal of Physical Therapy Science*, 21(1), 75-80. doi:10.1589/jpts.21.75
- Kim, H. D., Han, J. T., & Cho, Y. H. (2009). The effectiveness of community-based Tai Chi training on balance control during stair descent by older adults. *Journal of Physical Therapy Science*, 21(4), 317-323.
- Logghe, I. H., Verhagen, A. P., Rademaker, A. C., Bierma-Zeestra, S. M., van Rossum, E., Faber, M. J., & Koes, B. W. (2010). The effects of Tai Chi on fall prevention, fear of falling and balance in older people: a meta-analysis. *Preventive medicine*, 51(3-4), 222-227.
- Mao, D. W., Li, J. X., & Hong, Y. (2006a). Plantar pressure distribution during Tai Chi exercise. *Arch Phys Med Rehabil*, 87(6), 814-820.
- Mao, D. W., Li, J. X., & Hong, Y. (2006b). The duration and plantar pressure distribution during one-leg stance in Tai Chi exercise. *Clinical biomechanics (Bristol, Avon)*, 21(6), 640-645.
- Mian, O. S., Narici, M. V., Minetti, A. E., & Baltzopoulos, V. (2007). Centre of mass motion during stair negotiation in young and older men. *Gait and Posture*, 26(3), 463-469.
- Polcyn, A. F., Lipsitz, L. A., Kemigan, D. C., & Collins, J. J. (1998). Age-related changes in the initiation of gait: degradation of central mechanisms for momentum generation. *Arch Phys Med Rehabil*, 79(12), 1582-1589.
- Reid, S. M., Novak, A. C., Brouwer, B., & Costigan, P. A. (2011). Relationship between stair ambulation with and without a handrail and centre of pressure velocities during stair ascent and descent. *Gait and Posture*, 34(4), 529-532.
- Scarborough, M. D., Krebs, D. E., & Harris, B. A. (1999). Quadriceps muscle strength and dynamic stability in elderly persons. *Gait and Posture*, 10(1), 10-20.
- Sheehan, R. C., & Gottschall, J. S. (2011). Stair walking transitions are an anticipation of the next stride. *Journal of electromyography and kinesiology*, 21(3), 533-541.
- Svanstrom, L. (1974). Falls on stairs: an epidemiological accident study. *Scand J Soc Med*, 2(3), 113-120.
- van Dieën, J. H., Sparjaard, M., Konemann, R., Bron, L., & Pijnappels, M. (2007). Balance control in stepping down expected and unexpected level changes. *Journal of biomechanics*, 40(16), 3641-3649.
- Winter, D. A., Patla, A. E., Ishac, M., & Gage, W. H. (2003). Motor mechanisms of balance during quiet standing. *Journal of electromyography and kinesiology*, 13(1), 49-56.
- Wu, G. (2008). Age-related differences in Tai Chi gait kinematics and leg muscle electromyography: a pilot study. *Arch Phys Med Rehabil*, 89(2), 351-357.

ACKNOWLEDGEMENTS: This study is supported by "Aim for the Top University Plan" of the National Taiwan Normal University and the Ministry of Education, Taiwan, R.O.C

17. Hughes CP, Berg L, Danziger WL, Cohen LA, Martin RL. A new clinical scale for the staging of dementia. *Br J Psychiatry*. 1982;140:566–572. <https://doi.org/10.1192/bjp.140.6.566>
18. Kim J, Shin W. How to Do Random Allocation (Randomization). *Clinics in Orthopedic Surgery*. 2014;6(1):103. <https://doi.org/10.4055/cios.2014.6.1.103>
19. Tinetti ME. Performance-oriented assessment of mobility problems in elderly patients. *J Am Geriatr Soc*. 1986;34(2):119–126. <https://doi.org/10.1111/j.1532-5415.1986.tb05480.x>
20. Ajrcan D. Validity and reliability of Turkish version of Tinetti balance and gait assessment. Pamukkale University Institute of Health Sciences; 2009.
21. Washburn RA, Smith KW, Jette AM, Janney CA. The physical activity scale for the elderly (PASE): development and evaluation. *J Clin Epidemiol*. 1992;46(2):153–162.
22. Ayvaz E, Kilinc M, Kirdi N. The Turkish version of the physical activity scale for the elderly (PASE): Its cultural adaptation, validation, and reliability. *Turk J Med Sci*. 2017;47(3):908–915. <https://doi.org/10.3906/tjpm-1605-7>
23. Hudes K. The Tampa scale of kinesiophobia and neck pain, disability, and range of motion: a narrative review of the literature. *J Clin Chiropr Assoc*. 2011;55(3):222–232.
24. Yilmaz OF, Yakut Y, Uygur F, Uluğ N. Turkish version of the Tampa scale for kinesiophobia and its test-retest reliability. *Physiother Rehabil*. 2011;22(1):44–49.
25. Clemson L, Cumming RG, Kendig H, Swann M, Heard R, Taylor K. The effectiveness of a community-based program for reducing the incidence of falls in the elderly: a randomised trial. *J Am Geriatr Soc*. 2004;52(9):1497–1499. <https://doi.org/10.1111/j.1532-5415.2004.52411.x>
26. Ekşi Uyruş P, Nahoşvan N. Reliability and validity of fall behavioral scale for older people. *Florence Nightingale J Nurs*. 2013;21(1):22–32.
27. Huang P, Fang B, Li BY, Chen SD. Exercise-related changes of networks in aging and mild cognitive impairment brain. *Front Aging Neurosci*. 2016;8:47. <https://doi.org/10.3389/fnagi.2016.00047>
28. Osoba MY, Rao AK, Agrawal SK, Lalwani AK. Balance and gait in the elderly: a contemporary review. *Laryngoscope Invest Otolaryngol*. 2019;4(1):143–153. <https://doi.org/10.1002/lio2.222>
29. Pirker W, Kalzarschlager R. Gait disorders in adults and the elderly: a clinical guide. *Wien Klin Wochenschr*. 2017;129(3):481–95. <https://doi.org/10.1007/s00508-016-1096-4>
30. Jia RX, Liang JH, Xu Y, Wang YQ. Effects of physical activity and exercise on the cognitive function of patients with Alzheimer disease: a meta-analysis. *BMC Geriatr*. 2019;19(1):181. <https://doi.org/10.1186/s12877-019-1175-2>
31. Barnes JN. Exercise, cognitive function, and aging. *Adv Physiol Educ*. 2015;39(2):55–62. <https://doi.org/10.1152/advan.00101.2014>
32. Du Z, Li Y, Li J, Zhou C, Li F, Yang X. Physical activity can improve cognition in patients with Alzheimer's disease: a systematic review and meta-analysis of randomized controlled trials. *Clin Interv Aging*. 2018;13:1593–1603. <https://doi.org/10.2147/CIA.S169565>
33. Liu HH, Yeh NC, Wu YF, Yang YR, Wang RY, Cheng FY. Effects of Tai Chi exercise on reducing falls and improving balance performance in Parkinson's disease: a meta-analysis. *Parkinsons Dis*. 2019;2019:9626934–9626938. <https://doi.org/10.1155/2019/9626934>
34. Zou L, Wang C, Tian Z, Wang H, Shu Y. Effect of Yang-style Tai Chi on gait parameters and musculoskeletal flexibility in healthy Chinese older women. *Sports*. 2017;5(5):52. <https://doi.org/10.3390/sports5030052>
35. Lesinski M, Hortobagyi T, Muehlbauer T, Gollhofer A, Granacher U. Effects of balance training on balance performance in healthy older adults: a systematic review and meta-analysis. *Sports Med*. 2015;45(12):1721–1730. <https://doi.org/10.1007/s00279-015-0325-y>
36. Low DC, Walsh GS, Arakstisji M. Effectiveness of exercise interventions to improve postural control in older adults: a systematic review and meta-analysis of centre of pressure measurements. *Sports Med*. 2017;47(1):101–112. <https://doi.org/10.1007/s00279-016-0359-0>
37. Sangsri B, Kordi M, Barilushemi F, Nabavi SM, Khaladadch S, Dasgupta M. A combined exercise model for improving muscle strength, balance, walking distance, and motor agility in multiple sclerosis patients: a randomized clinical trial. *Iran J Neurosci*. 2016;15(3):111–120.
38. Gunn AH, Schwartz TA, Arbeeve LS, et al. Fear of movement and associated factors among adults with symptomatic knee osteoarthritis. *Arthritis Care Res*. 2017;69(12):1826–1833. <https://doi.org/10.1002/acr.23276>

How to cite this article: Birimoglu Okuyan C, Deveci E. The effectiveness of Tai Chi Chuan on fear of movement, prevention of falls, physical activity, and cognitive status in older adults with mild cognitive impairment: A randomized controlled trial. *Perspect Psychiatr Care*. 2020;1–9. <https://doi.org/10.1111/ppc.12684>

- Prieto, T.E., Myklebust, J.B., Hoffmann, R.G., & Lovett, E.G. (1996). Measures of postural steadiness: Differences between healthy young and elderly adults. *IEEE Transactions on Biomedical Engineering*, 43(9), 956–966. PubMed doi:10.1109/10.532130
- Scianni, A. (2015). Tai Chi improves balance and prevents falls in people with Parkinson's disease. *Journal of Physiotherapy*, 61(1), 44. PubMed doi:10.1016/j.jphys.2014.11.008
- Song, Q., Tian, X., Wong, D., Zhang, C., Sun, W., Cheng, P., & Mao, D. (2017). Effects of Tai Chi Exercise on body stability among the elderly during stair descent under different levels of illumination. *Research in Sports Medicine*, 25(2), 197–208. PubMed doi:10.1080/15438627.2017.1282363
- Sturnieks, D.L., George, R.S., & Lord, S.R. (2008). Balance disorders in the elderly. *Clinical Neurophysiology*, 38(6), 467–478. doi:10.1016/j.neucli.2008.09.001
- Sun, W., Zhang, C., Song, Q., Li, W., Cong, Y., Chang, S., ... Hong, Y. (2016). Effect of 1-year regular Tai Chi on neuromuscular reaction in elderly women: A randomized controlled study. *Research in Sports Medicine*, 24(2), 145–156. doi:10.1080/15438627.2015.1126280
- Tanaka, T., Takeda, H., Izumi, T., Ino, S., & Ifukube, T. (1999). Effects on the location of the centre of gravity and the foot pressure contribution to standing balance associated with ageing. *Ergonomics*, 42(7), 997–1010. PubMed doi:10.1080/001401399185261
- Tsang, W.W.N. (2013). Tai Chi training is effective in reducing balance impairments and falls in patients with Parkinson's disease. *Journal of Physiotherapy*, 59(1), 55. PubMed doi:10.1016/S1836-9553(13)70148-6
- Wang, S.J., Xu, D.Q., & Li, J.X. (2017). Effects of regular Tai Chi practice and jogging on neuromuscular reaction during lateral postural control in older people. *Research in Sports Medicine*, 25(1), 111–117. PubMed doi:10.1080/15438627.2016.1258649
- Won, S.H., Kim, J.C., & Oh, D.W. (2015). Effects of a novel walking training program with postural correction and visual feedback on walking function in patients with post-stroke hemiparesis. *Journal of Physical Therapy Science*, 27(8), 2581–2583. PubMed doi:10.1589/jpts.27.2581
- Xu, D.Q., Hong, Y., & Li, J.X. (2008). Tai Chi exercise and muscle strength and endurance in older people. In Y. Hong (Ed.), *Tai Chi Chuan: State of the art international research* (pp. 20–29). Basel, Switzerland: Karger.
- Yümin, E.T., Şimşek, T.T., Sertel, M., & Ankaralı, H. (2016). The effect of age and body mass index on plantar cutaneous sensation in healthy women. *Journal of Physical Therapy Science*, 28(9), 2587–2595. doi:10.1589/jpts.28.2587
- Zhang, C., Sun, W., Yu, B., Song, Q., & Mao, D. (2014). Effects of exercise on ankle proprioception in adult women during 16 weeks of training and eight weeks of detraining. *Research in Sports Medicine*, 23(1), 1–12.
- Zhou, J., Chang, S., Yan, C., Qin, M., Sun, W., Lian, J., ... Hong, Y. (2015). Effects of 24 weeks of Tai Chi exercise on postural control among elderly women. *Research in Sports Medicine*, 23(3), 302–314. PubMed doi:10.1080/15438627.2015.1040918

- [9] Hong, Y.L., Li, J.X., Robinson, P. (2015). Balance control, flexibility and cardiorespiratory fitness among older Tai Chi practitioners. *Br J Sports Med*, 34 (2000), pp. 29-34.
- [10] Kirkwood, R.N.; Gomes, H.D.; Sampaio, R.F.; Culham, E.; Costigan, P. (2007) Biomechanical analysis of hip and knee joints during gait in elderly subjects. *Acta Orthop. Bras.* 2007, 15, 267-271.
- [11] König, P. R., Galarza, E., Goulart, N., et al (2014). Effects of Tai Chi Chuan on the elderly balance: a semi-experimental study. *Revista Brasileira de Geriatria e Gerontologia Rev. bras. geriatr. gerontol.* vol.17 no.2 Rio de Janeiro 2014.
- [12] Kwon, J.W.; Son, S.M.; Lee, N.K. (2015) Changes of kinematic parameters of lower extremities with gait speed: A 3D motion analysis study. *J. Phys. Ther. Sci.* 27, 477-479.
- [13] Lamoureux, E.; Sparrow, W.; Murphy, A.; Newton, R. (2013) The effects of improved strength on obstacle negotiation in community-living older adults. *Gait Posture*, 17, 273-283.
- [14] Lau, C., Lai, J.S., Wong, M.K., Yu, M.L. (2012). Cardiorespiratory function, flexibility and body composition among geriatric Tai Chi Chuan Practitioners. *Arch Phys Med Rehabil*, 77 (1996), pp. 612-616.
- [15] Lelard, T., Doutrelot, P.L., David, P., Ahmaidi, S. (2010). Effects of a 12-week Tai Chi Chuan program versus a balance training program on postural control and walking ability in older people. *Arch Phys Med Rehabil*, 91 (2010), pp. 9-14.
- [16] Li, F., Harret, P. Fisher, E., McAuley, E. (2004). Tai Chi: improving functional balance and predicting subsequent falls in older persons. *Med Sci Sports Exerc*, 36 (2004), pp. 2046-2052.
- [17] Maki, B.E (2007). Gait changes in older adults: Predictors of falls or indicators of fear?. *J. Am. Geriatr. Soc.*, 45, 313-320.
- [18] Miyamoto, S.T., Lombardi, J.I., Berg, K.O., Natour J., Ramos, L.R. (2004). Brazilian Version of Berg Balance Scale. *Braz J Med Biol Res* 2004;37(9):1411-21.
- [19] Silvana, S., Santos, I, Da Silva, M., Barbosa, L., Porto, D., Pelzer, M., Silva, R. (2012). Risk of falls in the elderly: an integrative review based on the North American Nursing Diagnosis Association. *Revista da Escola de Enfermagem da USP*. vol.46 no.5 São Paulo Oct. 2012
- [20] Taylor, D., Hale, L., Schluter, P., et al. (2007). Effectiveness of tai chi as a community-based falls prevention intervention: a randomized controlled trial. *J Am Geriatr Soc*, 5 (2012), pp. 841-848.
- [21] Wayne, P.M., Kapchuk, T.J. (2015). Challenges inherent to tai chi research: part I – Tai Chi as a complex multicomponent intervention. *J Altern Complement Med*, 14 (2008), pp. 95-102.
- [22] Wayne, P.M.; Hausdorff, J.M.; Lough, M.; Gow, B.; Lipsitz, L.; Novak, V.; Macklin, E.; Peng, C.; Manor, B. (2015). Tai chi training may reduce dual task gait variability, a potential mediator of fall risk, in healthy older adults: Cross-sectional and randomized trial studies. *Front. Hum. Neurosci.* 9, 332.
- [23] Yeh, G., Wang, C., Wayne, P.M., Phillips, R.S. (2008). The effect of tai chi exercise on blood pressure: a systematic review. *Preventive Cardiology*. 2008 Spring;11(2):82-9.
- [24] The health benefits of tai chi. *Harvard Health Publications, Harvard Medical School*. Retrieved from <http://www.health.harvard.edu/staying-healthy/the-health-benefits-of-tai-chi>
- [25] 2012 Tai Chi named as perfect exercise for the elderly. *The Telegraph*. Retrieved from <http://www.telegraph.co.uk/news/health/elder/9187133/Tai-Chi-named-as-perfect-exercise-for-the-elderly.html>
- [26] Tai Chi as an intervention to reduce falls and improve balance function in the elderly: A meta-analysis of randomized controlled trials. Retrieved from <https://doi.org/10.1016/j.cure.2015.10.003>

